

# Note: Governing Equations of General 3D duct flow

Email: jiaqi\_wang@sjtu.edu.cn

## I. Hard-walled cylindrical ducts as basis function

### A. Infinite straight duct mode

We began from the Helmholtz equation:

$$\nabla^2 \psi = \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial \psi}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 \psi}{\partial \theta^2} = -\alpha^2 \psi \quad (1)$$

Using separation of variables, Circular symmetry: modes have the form :  $\psi = F(r)G(\theta)$ ,

Then we have:

$$\begin{aligned} \left( \frac{\partial^2 F}{\partial r^2} + \frac{1}{r} \frac{\partial F}{\partial r} \right) G + \frac{F}{r^2} \frac{\partial^2 G}{\partial \theta^2} &= -\alpha^2 FG \\ \text{Then,} \quad \left( \frac{\partial^2 F}{\partial r^2} + \frac{1}{r} \frac{\partial F}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 G}{\partial \theta^2} &= -\alpha^2 \end{aligned} \quad (2)$$

We assume that:

Due to periodicity, we require that  $\Phi$  satisfy,

$$\frac{d^2 G}{d\theta^2} = -m^2 G \rightarrow \Phi(\theta) = e^{\pm im\theta} \quad (3)$$

Thus, we have

$$F'' + \frac{1}{r} F' + \left( \alpha^2 - \frac{m^2}{r^2} \right) = 0 \rightarrow F(r) = J_m(\alpha r) \quad (4)$$

Circular symmetry  $\psi = F(r)G(\theta)$ : modes explicitly given by:

$$\psi = J_m(\alpha_{m\mu} r) e^{\pm im\theta} \quad (5)$$

Hard walls:

$$J'_m(\alpha R) = 0 \rightarrow \alpha_{m\mu} = \frac{j'_{m\mu}}{R} \quad (6)$$

Soft walls without flow:

$$Z\alpha_{m\mu}J'_m(\alpha_{m\mu}R) = -iw\rho_0 J_m(\alpha_{m\mu}R) \rightarrow \alpha_{m\mu}(Z) \quad (7)$$

Soft walls with flow:

$$Z\alpha_{m\mu}J'_m(\alpha_{m\mu}R) = (w - U_0\kappa_{m\mu})J_m(\alpha_{m\mu}R) \rightarrow \alpha_{m\mu}(Z) \quad (8)$$

A complete solution may be written as:

$$p(x, r, \theta) = \sum_{m=-\infty}^{\infty} \sum_{\mu=1}^{\infty} (A_{m\mu}e^{-i\kappa_{m\mu}x} + B_{m\mu}e^{i\kappa_{m\mu}x})U_{m\mu}(r)e^{im\phi} \quad (9)$$

In a hard-walled duct  $U_{m\mu}e^{-im\theta}$  are orthogonal. Normalise such that:

$$\int_0^{2\pi} \int_0^R U_{m\mu}(r)e^{-im\theta} U_{nv}(r)e^{-in\theta} r dr = 2\pi\delta_{\mu\nu}\delta_{mn} \quad (10)$$

Source expansion If  $p(0, t, \theta) = p_0(r, \theta)$  is source in hard-walled duct, then for  $x > 0$

$$\begin{aligned} p_0(r, \theta) &= \sum_{m=-\infty}^{\infty} \sum_{\mu=1}^{\infty} (A_{m\mu}e^{-i\kappa_{m\mu}0})U_{m\mu}(r)e^{-im\theta} \\ p_0(r, \theta) \underline{U_{nv}(r)e^{-in\theta} r} &= \sum_{m=-\infty}^{\infty} \sum_{\mu=1}^{\infty} (A_{m\mu}e^{-i\kappa_{m\mu}0})U_{m\mu}(r)e^{-im\theta} \underline{U_{nv}(r)e^{-in\theta} r} \\ \int_0^{2\pi} \int_0^R p_0(r, \theta) \underline{U_{nv}(r)e^{-in\theta} r} dr d\theta &= \int_0^{2\pi} \int_0^R \sum_{m=-\infty}^{\infty} \sum_{\mu=1}^{\infty} (A_{m\mu}e^{-i\kappa_{m\mu}0})U_{m\mu}(r)e^{-im\theta} \underline{U_{nv}(r)e^{-in\theta} r} dr d\theta \\ \int_0^{2\pi} \int_0^R p_0(r, \theta) \underline{U_{nv}(r)e^{-in\theta} r} dr d\theta &= \sum_{m=-\infty}^{\infty} \sum_{\mu=1}^{\infty} A_{m\mu} \int_0^{2\pi} \int_0^R U_{m\mu}(r)e^{-im\theta} \underline{U_{nv}(r)e^{-in\theta} r} dr d\theta \\ A_{nv} &= \frac{1}{2\pi} \int_0^{2\pi} \int_0^R p_0(r, \theta) \underline{U_{nv}(r)e^{-in\theta} r} dr d\theta \end{aligned} \quad (11)$$

and  $B_{nv} = 0$ . The same for  $x < 0$  with  $A_{nv}$  and  $B_{nv}$  interchanged.

A finite number of modes (cut-on modes) survive at large distances. Just 1 mode if  $kR \ll 1$ : only  $A_{01}$  important.

## B. General duct mode

The pressure and velocity can now be expressed as Fourier series. Upper indices shall be used to denote temporal decompositions:

$$\begin{aligned}
\hat{p} &= \sum_{a=-\infty}^{\infty} P^a(\mathbf{x}) e^{-iawt} \\
\hat{u} &= \sum_{a=-\infty}^{\infty} U^a(\mathbf{x}) e^{-iawt} \\
\hat{v} &= \sum_{a=-\infty}^{\infty} V^a(\mathbf{x}) e^{-iawt} \\
\hat{w} &= \sum_{a=-\infty}^{\infty} W^a(\mathbf{x}) e^{-iawt}
\end{aligned} \tag{12}$$

$$\begin{aligned}
P^a &= \sum_{\alpha=0}^{\infty} P_{\alpha m \mu}^a(s) \psi_{\alpha m \mu}(s, r, \theta) = \sum_{m=-\infty}^{\infty} \sum_{\mu=1}^{\infty} P_{\alpha m \mu}^a(s) \psi_{\alpha m \mu}(s, r, \theta) \\
U^a &= \sum_{\alpha=0}^{\infty} U_{\alpha m \mu}^a(s) \psi_{\alpha m \mu}(s, r, \theta) = \sum_{m=-\infty}^{\infty} \sum_{\mu=1}^{\infty} U_{\alpha m \mu}^a(s) \psi_{\alpha m \mu}(s, r, \theta) \\
V^a &= \sum_{\alpha=0}^{\infty} V_{\alpha m \mu}^a(s) \psi_{\alpha m \mu}(s, r, \theta) = \sum_{m=-\infty}^{\infty} \sum_{\mu=1}^{\infty} V_{\alpha m \mu}^a(s) \psi_{\alpha m \mu}(s, r, \theta) \\
W^a &= \sum_{\alpha=0}^{\infty} W_{\alpha m \mu}^a(s) \psi_{\alpha m \mu}(s, r, \theta) = \sum_{m=-\infty}^{\infty} \sum_{\mu=1}^{\infty} W_{\alpha m \mu}^a(s) \psi_{\alpha m \mu}(s, r, \theta)
\end{aligned} \tag{13}$$

### 1. Discussion of property of $\psi$

In Jams's thesis,  $\psi_{\alpha} = C_{\alpha m \mu} J_m(\frac{j'_{m \mu} r}{h}) \cos(m\phi - \xi\pi/2)$ . As he mentioned, it was merely a matter of preference - I find it easier to visualise the modes as being "symmetric" and "anti-symmetric" along the plane of torsion free ducts, but the other method is equally as valid.

However, there still a problem, which may cause error, but is not mentioned in the thesis:

$$\int_0^{2\pi} \cos(m\phi - \xi\pi/2)^2 d\theta = \frac{1 + \cos(2m\phi - \xi\pi)}{2} |_{0}^{2\pi} = \begin{cases} \pi, m \neq 0 \\ 2\pi, m = 0, \xi = 0 \\ 0, m = 0, \xi = 1 \end{cases} \tag{14}$$

In the note, we try to introduce the common solution of  $\psi$  may have the form the same as the hard walls modes:

$$\psi_{m \mu}(r) = C_{\alpha m \mu} J_m(\frac{j'_{m \mu} r}{h}) e^{im\phi} \tag{15}$$

where may be normalized according to:

$$\int_0^{2\pi} \int_0^h \psi_{\alpha m \mu} \psi_{\beta n \nu} r dr d\theta = \delta_{\mu\nu} \delta_{mn} \tag{16}$$

In fact, we know that:

$$\begin{aligned} \int_0^{2\pi} e^{im\phi} e^{im\phi} d\theta &= 0 \\ \int_0^{2\pi} e^{im\phi} e^{-im\theta} d\theta &= 2\pi \end{aligned} \quad (17)$$

The orthogonality relation of Bessel function, with  $J_{-m}(z) = (-1)^m J_m(z)$

$$\begin{aligned} \int_0^h r J_m\left(\frac{j'_{m\mu} r}{h}\right) J_m\left(\frac{j'_{mv} r}{h}\right) dr &= 0, \mu \neq v \\ \int_0^h r J_m\left(\frac{j'_{m\mu} r}{h}\right) J_{-m}\left(\frac{j'_{-mv} r}{h}\right) dr \\ &= (-1)^m \int_0^h r J_m\left(\frac{j'_{m\mu} r}{h}\right) J_m\left(\frac{j'_{mv} r}{h}\right) dr = 0, \mu \neq v \end{aligned} \quad (18)$$

That changes our idea of normalization to:

$$\underline{\underline{\int_0^{2\pi} \int_0^h \psi_{\alpha_{m\mu}} \psi_{\beta_{nv}} r dr d\theta = (-1)^m \delta_{\mu\nu} \delta_{m,-n}}} \quad (19)$$

### C. Normalised Modes $\rightarrow C_{\alpha_{m\mu}}$

Relation involving intergrals:

$$\begin{aligned} &\frac{2 \int \alpha^2 x J_m(\alpha x)^2 dx = (\alpha^2 x^2 - m^2) J_m(\alpha x)^2 + \alpha^2 x^2 J'_m(\alpha x)^2}{\int_0^h \alpha^2 x J_m(\alpha x)^2 dx = [(\alpha^2 x^2 - m^2) J_m(\alpha x)^2 + \alpha^2 x^2 J'_m(\alpha x)^2]|_0^h} \\ &= [(\alpha^2 x^2 - m^2) J_m(\alpha x)^2 + \alpha^2 x^2 J'_m(\alpha x)^2]|_h - [(\alpha^2 x^2 - m^2) J_m(\alpha x)^2 + \alpha^2 x^2 J'_m(\alpha x)^2]|_0 \\ &= [(\alpha^2 h^2 - m^2) J_m(\alpha h)^2 + \alpha^2 h^2 J'_m(\alpha h)^2] - [(\alpha^2 0^2 - m^2) J_m(\alpha 0)^2 + \alpha^2 0^2 J'_m(\alpha 0)^2] \\ &= (\alpha^2 h^2 - m^2) J_m(\alpha h)^2 + \alpha^2 h^2 J'_m(\alpha h)^2 \end{aligned} \quad (20)$$

With hard-walled boudary condition:

$$J'_m(\alpha h) = 0 \rightarrow \alpha_{m\mu} = \frac{j'_{m\mu}}{h} \text{ (eigenvalues)} \quad (21)$$

Then, we have(ref: Rienstra-Fundamentals of Duct Acoustics-(55)):

$$\begin{aligned} \int_0^h r J_m(\alpha r) J_{-m}(\alpha r) dr &= (-1)^m \int_0^h r J_m(\alpha r)^2 dr \\ &= (-1)^m \frac{1}{2\alpha_{m\mu}^2} (\alpha_{m\mu}^2 h^2 - m^2) J_m(\alpha_{m\mu} h)^2 \\ &= (-1)^m \left( \frac{J_m(\alpha_{m\mu} h) \sqrt{(h^2 - \frac{m^2}{\alpha_{m\mu}^2})}}{\sqrt{2}} \right)^2 \\ &= (-1)^m \left( \frac{h^2}{2} \left( 1 - \frac{m^2}{j'^2_{m\mu}} \right) J_m^2(j'_{m\mu}) \right) \end{aligned} \quad (22)$$

Thus, with  $\int_0^{2\pi} e^{im\phi} e^{-im\theta} d\theta = 2\pi$ ,  $\psi_{\alpha_{m\mu}}(r) = C_{\alpha_{m\mu}} J_m(\frac{j'_{m\mu} r}{h}) e^{im\phi}$ , we have:

$$C_{\alpha_{m\mu}} = \frac{i^m}{\sqrt{(\pi h^2(1 - \frac{m^2}{j'^2_{m\mu}})J_m^2(j'_{m\mu}))}} \quad (23)$$

except for :  $C_{\alpha_{01}} = \frac{1}{\sqrt{\pi}h}$

for  $\int_0^{2\pi} \int_0^h \psi_{\alpha_{m\mu}} \psi_{\beta_{n\nu}} r dr d\theta = \delta_{\mu\nu} \delta_{m,-n} = \hat{\delta}_{\alpha\beta} = I$ .

#### D. Slowly varying ducts

waiting for updating.....

#### E. Orthogonal-eigenvector

ref:<https://www.mathworks.com/help/matlab/ref/eigs.html>

Eigenvectors, returned as a matrix. The columns in V correspond to the eigenvalues along the diagonal of D. The form and normalization of V depends on the combination of input arguments:  
 $[V,D] = \text{eigs}(A)$  returns matrix V, whose columns are the eigenvectors of A such that  $A^*V = V^*D$ . The eigenvectors in V are normalized so that the 2-norm of each is 1.

If A is symmetric, then the eigenvectors, V, are orthonormal.

$[V,D] = \text{eigs}(A,B)$  returns V as a matrix whose columns are the generalized eigenvectors that satisfy  $A^*V = B^*V^*D$ . The 2-norm of each eigenvector is not necessarily 1.

If B is symmetric positive definite, then the eigenvectors in V are normalized so that the B-norm of each is 1. If A is also symmetric, then the eigenvectors are B-orthonormal.

We could further study this question!!

if we can use the GramSchmidt mode as basis??

## II. Mass equation

Mass conservation:

$$-ia\kappa P^a + \nabla \cdot \mathbf{U}^a = \sum_{b=-\infty}^{+\infty} (-P^{a-b} \nabla \cdot \mathbf{U}^b - \mathbf{U}^{a-b} \cdot \nabla P^b - \frac{B}{2A} i a k P^b P^{a-b}) \quad (24)$$

First, derivation of eq1:

We know that:

$$h_s = 1 - \kappa r \cos(\phi), h_r = 1, h_\theta = r \quad (25)$$

Then,

$$\begin{aligned} \nabla \cdot \mathbf{U}^a &= \frac{1}{h_1 h_2 h_3} \left[ \frac{\partial(v_1 h_2 h_3)}{\partial w_1} + \frac{\partial(v_2 h_3 h_1)}{\partial w_2} + \frac{\partial(v_3 h_1 h_2)}{\partial w_3} \right] \\ &= \frac{1}{r(1 - \kappa r \cos(\phi))} \left[ \frac{\partial(U^a r)}{\partial s} + \frac{\partial(V^a r(1 - \kappa r \cos(\phi)))}{\partial r} + \frac{\partial(W^a(1 - \kappa r \cos(\phi)))}{\partial \theta} \right] \end{aligned} \quad (26)$$

Thus, we have the mass equation, approximate RHS by:

$$\begin{aligned} \nabla \cdot \mathbf{U}^b &= ib\kappa P^b + o(M^2) \\ \nabla P^b &= ib\kappa \mathbf{U}^b + o(M^2) \end{aligned} \quad (27)$$

Then we have

$$\begin{aligned} -ia\kappa P^a + \frac{1}{r(1 - \kappa r \cos(\phi))} \left[ \frac{\partial(U^a r)}{\partial s} + \frac{\partial(V^a r(1 - \kappa r \cos(\phi)))}{\partial r} + \frac{\partial(W^a(1 - \kappa r \cos(\phi)))}{\partial \theta} \right] \\ = \sum_{b=-\infty}^{+\infty} (-ib\kappa P^{a-b} P^b - ib\kappa U^{a-b} U^b - ib\kappa V^{a-b} V^b - ib\kappa W^{a-b} W^b - \frac{B}{2A} iak P^b P^{a-b}) \end{aligned} \quad (28)$$

The fourier harmonics are expanded as follows:

$$\begin{aligned} P^a &= \sum_{\beta=0}^{\infty} P_{\beta}^a(s) \psi_{\beta}(s, r, \theta) \\ U^a &= \sum_{\beta=0}^{\infty} U_{\beta}^a(s) \psi_{\beta}(s, r, \theta) \\ V^a &= \sum_{\beta=0}^{\infty} V_{\beta}^a(s) \psi_{\beta}(s, r, \theta) \\ W^a &= \sum_{\beta=0}^{\infty} W_{\beta}^a(s) \psi_{\beta}(s, r, \theta) \end{aligned} \quad (29)$$

with normalized relation:

$$\int_0^{2\pi} \int_0^h \psi_{\alpha} \psi_{\beta} r dr d\theta = \hat{\delta}_{\alpha\beta} \quad (30)$$

Reorganize the eq5:

$$\begin{aligned} -ia\kappa P^a(1 - \kappa r \cos(\phi)) + \frac{1}{r} \frac{\partial(U^a r)}{\partial s} + \frac{1}{r} \frac{\partial(V^a r(1 - \kappa r \cos(\phi)))}{\partial r} + \frac{1}{r} \frac{\partial(W^a(1 - \kappa r \cos(\phi)))}{\partial \theta} \\ = (1 - \kappa r \cos(\phi)) \sum_{b=-\infty}^{+\infty} (-ib\kappa P^{a-b} P^b - ib\kappa U^{a-b} U^b - ib\kappa V^{a-b} V^b - ib\kappa W^{a-b} W^b - \frac{B}{2A} iak P^b P^{a-b}) \end{aligned} \quad (31)$$

Intergal and insert eq 6, 7 into eq 8:

1. the first term:

$$\begin{aligned}
& \int_0^{2\pi} \int_0^h \psi_\alpha r [-ia\kappa(1 - \kappa \cos(\phi)) P^a] dr d\theta \\
&= -ia\kappa \sum_{\beta=0}^{\infty} \int_0^{2\pi} \int_0^h \psi_\alpha \psi_\beta r [(1 - \kappa \cos(\phi))] dr d\theta P_\beta^a \\
&\Rightarrow -ia\kappa \Psi_{\alpha\beta} [r(1 - \kappa \cos(\phi))] P_\beta^a \\
&\quad (\text{summation convention})
\end{aligned} \tag{32}$$

2. the second term:

From 1.1 as example, we know that

$$\begin{aligned}
& \int_0^{2\pi} \int_0^h \frac{\partial}{\partial s} [r U^{a-b} U^b \psi_\alpha] dr d\theta \\
&= \frac{\partial}{\partial s} \int_0^{2\pi} \int_0^h [r U^{a-b} U^b \psi_\alpha] dr d\theta - \int_0^{2\pi} \frac{dh(s)}{ds} [r U^{a-b} U^b \psi_\alpha]_{r=h} d\theta
\end{aligned} \tag{33}$$

$$\frac{d}{d\alpha} \left( \int_{a(\alpha)}^{b(\beta)} f(x, \alpha) dx \right) - 0 + \frac{da(\alpha)}{d\alpha} f(a(\alpha), \alpha) = \int_{a(\alpha)}^{b(\beta)} \frac{\partial}{\partial \alpha} f(x, \alpha) dx \tag{34}$$

$$\begin{aligned}
& \int_0^{2\pi} \int_0^h \psi_\alpha r \left[ \frac{1}{r} \frac{\partial(U^a r)}{\partial s} \right] dr d\theta \\
&= \sum_{\beta=0}^{\infty} \int_0^{2\pi} \int_0^h \psi_\alpha r \frac{\partial(\psi_\beta U_\beta^a)}{\partial s} dr d\theta \\
&= \int_0^{2\pi} \int_0^h \psi_\alpha \psi_\beta r \frac{\partial(U_\beta^a)}{\partial s} dr d\theta + \int_0^{2\pi} \int_0^h \psi_\alpha r \frac{\partial(\psi_\beta)}{\partial s} dr d\theta U_\beta^a \\
&= \int_0^{2\pi} \int_0^h \frac{\partial \psi_\alpha \psi_\beta r U_\beta^a}{\partial s} dr d\theta - \int_0^{2\pi} \int_0^h \frac{\partial \psi_\alpha \psi_\beta r}{\partial s} U_\beta^a dr d\theta \\
&+ \int_0^{2\pi} \int_0^h r \frac{\partial(\psi_\alpha \psi_\beta)}{\partial s} dr d\theta U_\beta^a - \int_0^{2\pi} \int_0^h \psi_\beta r \frac{\partial(\psi_\alpha)}{\partial s} dr d\theta U_\beta^a \\
&= \sum_{\beta=0}^{\infty} \frac{\partial}{\partial s} \int_0^{2\pi} \int_0^h [\psi_\alpha \psi_\beta r U_\beta^a] dr d\theta - \sum_{\beta=0}^{\infty} \int_0^{2\pi} \frac{dh(s)}{ds} [\psi_\alpha \psi_\beta r U_\beta^a]_{r=h} d\theta \\
&\quad - \sum_{\beta=0}^{\infty} \int_0^{2\pi} \int_0^h \frac{\partial(\psi_\alpha)}{\partial s} \psi_\beta r dr d\theta U_\beta^a \\
&\Rightarrow \frac{d}{ds} (U_\beta^a \hat{\delta}_{\alpha\beta}) + 0(\text{periodic}) - \Psi_{\{\alpha\}\beta} [r] U_\beta^a
\end{aligned} \tag{35}$$

mark:0(periodic), which is similar in the follow momentum equation, but not eliminate in time.

3. the third term:

$$\begin{aligned}
& \int_0^{2\pi} \int_0^h \psi_\alpha r \left[ \frac{1}{r} \frac{\partial(V^a r(1 - \kappa r \cos(\phi)))}{\partial r} \right] dr d\theta \\
&= \sum_{\beta=0}^{\infty} \int_0^{2\pi} \int_0^h \psi_\alpha \frac{\partial(\psi_\beta r(1 - \kappa r \cos(\phi)))}{\partial r} dr d\theta V_\beta^a \\
&= \int_0^{2\pi} \int_0^h \frac{\partial(\psi_\alpha \psi_\beta r(1 - \kappa r \cos(\phi)))}{\partial r} dr d\theta V_\beta^a - \int_0^{2\pi} \int_0^h \psi_\beta r(1 - \kappa r \cos(\phi)) \frac{\partial(\psi_\alpha)}{\partial r} dr d\theta V_\beta^a \\
&= 0(\text{periodic}) - \Psi_{[\alpha]\beta}[r(1 - \kappa r \cos(\phi))] V_\beta^a
\end{aligned} \tag{36}$$

4. the fourth term:

$$\begin{aligned}
& \int_0^{2\pi} \int_0^h \psi_\alpha r \left[ \frac{1}{r} \frac{\partial(W^a(1 - \kappa r \cos(\phi)))}{\partial \theta} \right] dr d\theta \\
&= \sum_{\beta=0}^{\infty} \int_0^{2\pi} \int_0^h \psi_\alpha \left[ \frac{\partial(\psi_\beta(1 - \kappa r \cos(\phi)))}{\partial \theta} \right] dr d\theta W_\beta^a \\
&= \int_0^{2\pi} \int_0^h \left[ \frac{\partial(\psi_\alpha \psi_\beta(1 - \kappa r \cos(\phi)))}{\partial \theta} \right] dr d\theta W_\beta^a - \int_0^{2\pi} \int_0^h \psi_\beta(1 - \kappa r \cos(\phi)) \left[ \frac{\partial(\psi_\alpha)}{\partial \theta} \right] dr d\theta W_\beta^a \\
&= \int_0^h [\psi_\alpha \psi_\beta(1 - \kappa r \cos(\phi))]_0^{2\pi} dr W_\beta^a - \int_0^{2\pi} \int_0^h \psi_\beta(1 - \kappa r \cos(\phi)) \left[ \frac{\partial(\psi_\alpha)}{\partial \theta} \right] dr d\theta W_\beta^a \\
&= 0 - \Psi_{(\alpha)\beta}[(1 - \kappa r \cos(\phi))] W_\beta^a
\end{aligned} \tag{37}$$

5. the RHS term:

$$\begin{aligned}
& \int_0^{2\pi} \int_0^h \psi_\alpha r [(1 - \kappa r \cos(\phi))] \sum_{b=-\infty}^{+\infty} (-ib\kappa P^{a-b}P^b - ib\kappa U^{a-b}U^b - ib\kappa V^{a-b}V^b - ib\kappa W^{a-b}W^b - \frac{B}{2A} iak P^b P^{a-b}) dr d\theta \\
&= \sum_{\beta=0}^{\infty} \sum_{\gamma=0}^{\infty} \int_0^{2\pi} \int_0^h \psi_\alpha \psi_\beta \psi_\gamma r (1 - \kappa r \cos(\phi)) dr d\theta \\
&\quad \sum_{b=-\infty}^{+\infty} (-ib\kappa P_\beta^{a-b}P_\gamma^b - ib\kappa U_\beta^{a-b}U_\gamma^b - ib\kappa V_\beta^{a-b}V_\gamma^b - ib\kappa W_\beta^{a-b}W_\gamma^b - ia\kappa \frac{B}{2A} P_\beta^{a-b} P_\gamma^b) \\
&= \sum_{\beta=0}^{\infty} \sum_{\gamma=0}^{\infty} \Psi_{\alpha\beta\gamma}[r(1 - \kappa r \cos(\phi))] \sum_{b=-\infty}^{+\infty} (-ib\kappa P_\beta^{a-b}P_\gamma^b - ib\kappa U_\beta^{a-b}U_\gamma^b - ib\kappa V_\beta^{a-b}V_\gamma^b - ib\kappa W_\beta^{a-b}W_\gamma^b - ia\kappa \frac{B}{2A} P_\beta^{a-b} P_\gamma^b)
\end{aligned} \tag{38}$$

Finally, we obtain the mass equation in the form of eigenfunction, the idea is same as Galerkin method:

$$\begin{aligned}
& \frac{dU_\beta^a}{ds} \hat{\delta}_{\alpha\beta} - \Psi_{\{\alpha\}\beta}[r] U_\beta^a - ia\kappa \Psi_{\alpha\beta}[r(1 - \kappa \cos(\phi))] P_\beta^a - \Psi_{[\alpha]\beta}[r(1 - \kappa r \cos(\phi))] V_\beta^a - \Psi_{(\alpha)\beta}[(1 - \kappa r \cos(\phi))] W_\beta^a \\
&= \Psi_{\alpha\beta\gamma}[r(1 - \kappa r \cos(\phi))] \sum_{b=-\infty}^{+\infty} (-ib\kappa P_\beta^{a-b}P_\gamma^b - ib\kappa U_\beta^{a-b}U_\gamma^b - ib\kappa V_\beta^{a-b}V_\gamma^b - ib\kappa W_\beta^{a-b}W_\gamma^b - ia\kappa \frac{B}{2A} P_\beta^{a-b} P_\gamma^b)
\end{aligned} \tag{39}$$

### III. Momentum equation

Momentum conservation:

$$-ia\kappa \mathbf{U}^a + \nabla P^a = \sum_{b=-\infty}^{\infty} (-\mathbf{U}^{a-b} \cdot \nabla \mathbf{U}^b + P^{a-b} \nabla P^b) \quad (40)$$

First, we know that

$$\nabla P^a = \sum_i \frac{1}{h_i} \frac{\partial f}{\partial w_i} \hat{h}_i = \frac{1}{1 - \kappa r \cos \phi} \frac{\partial P^a}{\partial s} \hat{e}_s + \frac{\partial P^a}{\partial r} \hat{e}_r + \frac{1}{r} \frac{\partial P^a}{\partial \theta} \hat{e}_{\theta} \quad (41)$$

The RHS term is a bit complex, with the divergence of a vector  $\mathbf{U}$  with its gradient, with

First, we know that

$$(\mathbf{v} \cdot \nabla) \mathbf{v}^b = \begin{cases} term1 : \mathcal{D}v_1^b + \frac{v_2^b}{h_2 h_1} (v_1 \frac{\partial h_1}{\partial \xi_2} - v_2 \frac{\partial h_2}{\partial \xi_1}) + \frac{v_3^b}{h_3 h_1} (v_1 \frac{\partial h_1}{\partial \xi_3} - v_2 \frac{\partial h_3}{\partial \xi_1}) \\ term2 : \mathcal{D}v_2^b + \frac{v_3^b}{h_3 h_2} (v_2 \frac{\partial h_2}{\partial \xi_3} - v_3 \frac{\partial h_3}{\partial \xi_2}) + \frac{v_1^b}{h_1 h_2} (v_2 \frac{\partial h_2}{\partial \xi_1} - v_1 \frac{\partial h_1}{\partial \xi_2}) \\ term3 : \mathcal{D}v_3^b + \frac{v_1^b}{h_1 h_3} (v_3 \frac{\partial h_3}{\partial \xi_1} - v_1 \frac{\partial h_1}{\partial \xi_3}) + \frac{v_2^b}{h_2 h_3} (v_3 \frac{\partial h_3}{\partial \xi_2} - v_2 \frac{\partial h_2}{\partial \xi_3}) \end{cases} \quad (42)$$

Besides,

$$\mathcal{D} = \frac{v_1}{h_1} \frac{\partial}{\partial \xi_1} + \frac{v_2}{h_2} \frac{\partial}{\partial \xi_2} + \frac{v_3}{h_3} \frac{\partial}{\partial \xi_3} \quad (43)$$

Thus, we have:

$$\begin{aligned}
& -\mathbf{U}^{a-b} \cdot \nabla \mathbf{U}^b = \\
& -\sum_{b=-\infty}^{\infty} \left\{ \begin{array}{l} term\mathcal{D}1 : \frac{U^{a-b}}{1-\kappa r \cos \phi} \frac{\partial U^b}{\partial s} + V^{a-b} \frac{\partial U^b}{\partial r} + \frac{W^{a-b}}{r} \frac{\partial U^b}{\partial \theta} \\ term\mathcal{D}2 : \frac{U^{a-b}}{1-\kappa r \cos \phi} \frac{\partial V^b}{\partial s} + V^{a-b} \frac{\partial V^b}{\partial r} + \frac{W^{a-b}}{r} \frac{\partial V^b}{\partial \theta} \\ term\mathcal{D}3 : \frac{U^{a-b}}{1-\kappa r \cos \phi} \frac{\partial W^b}{\partial s} + V^{a-b} \frac{\partial W^b}{\partial r} + \frac{W^{a-b}}{r} \frac{\partial W^b}{\partial \theta} \end{array} \right. \\
& -\sum_{b=-\infty}^{\infty} \left\{ \begin{array}{l} term\mathcal{X}1 : \frac{V^b}{1-\kappa r \cos \phi} (U^{a-b} \frac{\partial(1-\kappa r \cos \phi)}{\partial r} - V^{a-b} \frac{\partial 1}{\partial s}) + \frac{W^b}{r(1-\kappa r \cos \phi)} (U^{a-b} \frac{\partial(1-\kappa r \cos \phi)}{\partial \theta} - V^{a-b} \frac{\partial r}{\partial s}) \\ term\mathcal{X}2 : \frac{W^b}{r} (V^{a-b} \frac{\partial 1}{\partial \theta} - W^{a-b} \frac{\partial r}{\partial r}) + \frac{U^b}{(1-\kappa r \cos \phi)1} (V^{a-b} \frac{\partial 1}{\partial s} - U^{a-b} \frac{\partial(1-\kappa r \cos \phi)}{\partial r}) \\ term\mathcal{X}3 : \frac{U^b}{(1-\kappa r \cos \phi)r} (W^{a-b} \frac{\partial h_3}{\partial s} - U^{a-b} \frac{\partial(1-\kappa r \cos \phi)}{\partial \theta}) + \frac{V^b}{1h_3} (W^{a-b} \frac{\partial h_3}{\partial r} - V^{a-b} \frac{\partial 1}{\partial \theta}) \end{array} \right. = \\
& \sum_{b=-\infty}^{\infty} \left\{ \begin{array}{l} term\mathcal{D}1 : -\frac{U^{a-b}}{1-\kappa r \cos \phi} \frac{\partial U^b}{\partial s} - V^{a-b} \frac{\partial U^b}{\partial r} - \frac{W^{a-b}}{r} \frac{\partial U^b}{\partial \theta} \\ term\mathcal{D}2 : -\frac{U^{a-b}}{1-\kappa r \cos \phi} \frac{\partial V^b}{\partial s} - V^{a-b} \frac{\partial V^b}{\partial r} - \frac{W^{a-b}}{r} \frac{\partial V^b}{\partial \theta} \\ term\mathcal{D}3 : -\frac{U^{a-b}}{1-\kappa r \cos \phi} \frac{\partial W^b}{\partial s} - V^{a-b} \frac{\partial W^b}{\partial r} - \frac{W^{a-b}}{r} \frac{\partial W^b}{\partial \theta} \end{array} \right. \\
& + \sum_{b=-\infty}^{\infty} \left\{ \begin{array}{l} term\mathcal{X}1 : \frac{\kappa \cos \phi}{1-\kappa r \cos \phi} U^{a-b} V^b - \frac{\kappa \sin \phi}{(1-\kappa r \cos \phi)} U^{a-b} W^b \\ term\mathcal{X}2 : \frac{W^{a-b} W^b}{r} - \frac{\kappa \cos \phi}{(1-\kappa r \cos \phi)} U^{a-b} U^b \\ term\mathcal{X}3 : \frac{\kappa \sin \phi}{(1-\kappa r \cos \phi)} U^{a-b} U^b - \frac{W^{a-b} V^b}{r} \end{array} \right. \quad (44)
\end{aligned}$$

Finally, we could derive the momentum conservation equation, with final term approximate by

eq 4:

$$\begin{aligned}
& \left\{ \begin{array}{l} -ia\kappa U^a + \frac{1}{1-\kappa r \cos \phi} \frac{\partial P^a}{\partial s} \\ -ia\kappa V^a + \frac{\partial P^a}{\partial r} \\ -ia\kappa W^a + \frac{1}{r} \frac{\partial P^a}{\partial \theta} \end{array} \right. \\
& = \sum_{b=-\infty}^{\infty} \left\{ \begin{array}{l} term\mathcal{D}1 : -\frac{U^{a-b}}{1-\kappa r \cos \phi} \frac{\partial U^b}{\partial s} - V^{a-b} \frac{\partial U^b}{\partial r} - \frac{W^{a-b}}{r} \frac{\partial U^b}{\partial \theta} \\ term\mathcal{D}2 : -\frac{U^{a-b}}{1-\kappa r \cos \phi} \frac{\partial V^b}{\partial s} - V^{a-b} \frac{\partial V^b}{\partial r} - \frac{W^{a-b}}{r} \frac{\partial V^b}{\partial \theta} \\ term\mathcal{D}3 : -\frac{U^{a-b}}{1-\kappa r \cos \phi} \frac{\partial W^b}{\partial s} - V^{a-b} \frac{\partial W^b}{\partial r} - \frac{W^{a-b}}{r} \frac{\partial W^b}{\partial \theta} \end{array} \right. \quad (45) \\
& + \sum_{b=-\infty}^{\infty} \left\{ \begin{array}{l} term\mathcal{X}1 : \frac{\kappa \cos \phi}{1-\kappa r \cos \phi} U^{a-b} V^b - \frac{\kappa \sin \phi}{(1-\kappa r \cos \phi)} U^{a-b} W^b \\ term\mathcal{X}2 : \frac{W^{a-b} W^b}{r} - \frac{\kappa \cos \phi}{(1-\kappa r \cos \phi)} U^{a-b} U^b \\ term\mathcal{X}3 : \frac{\kappa \sin \phi}{(1-\kappa r \cos \phi)} U^{a-b} U^b - \frac{W^{a-b} V^b}{r} \end{array} \right. + \sum_{b=-\infty}^{\infty} \left\{ \begin{array}{l} ib\kappa P^{a-b} U^b \\ ib\kappa P^{a-b} V^b \\ ib\kappa P^{a-b} W^b \end{array} \right.
\end{aligned}$$

Now, we are going to project on  $\psi$ , it may be a little complex, we will doing step by step.

### A. Momentum $e^s$ term

First, deal with the  $e^s$  term:

$$\begin{aligned}
& -ia\kappa U^a + \frac{1}{1 - \kappa r \cos\phi} \frac{\partial P^a}{\partial s} \\
= & \sum_{b=-\infty}^{\infty} \text{termD1} : -\frac{U^{a-b}}{1 - \kappa r \cos\phi} \frac{\partial U^b}{\partial s} - V^{a-b} \frac{\partial U^b}{\partial r} - \frac{W^{a-b}}{r} \frac{\partial U^b}{\partial \theta} \\
& + \sum_{b=-\infty}^{\infty} \text{termX1} : \frac{\kappa \cos\phi}{1 - \kappa r \cos\phi} U^{a-b} V^b - \frac{\kappa \sin\phi}{(1 - \kappa r \cos\phi)} U^{a-b} W^b \\
& + ib\kappa P^{a-b} U^b
\end{aligned} \tag{46}$$

Multiply  $(1 - \kappa r \cos\phi)$ , we have:

$$\begin{aligned}
& -ia\kappa(1 - \kappa r \cos\phi) U^a + \frac{\partial P^a}{\partial s} \\
= & \text{termD1} : \sum_{b=-\infty}^{\infty} -U^{a-b} \frac{\partial U^b}{\partial s} - (1 - \kappa r \cos\phi) V^{a-b} \frac{\partial U^b}{\partial r} - (1 - \kappa r \cos\phi) \frac{W^{a-b}}{r} \frac{\partial U^b}{\partial \theta} \\
& + \text{termX1} : \sum_{b=-\infty}^{\infty} \kappa \cos\phi U^{a-b} V^b - \kappa \sin\phi U^{a-b} W^b + \\
& \text{termP1} : \sum_{b=-\infty}^{\infty} ib\kappa(1 - \kappa r \cos\phi) P^{a-b} U^b
\end{aligned} \tag{47}$$

$\int \int X X r \psi_\alpha dr d\theta$ , we have:

$$RHS = \underbrace{\int_0^{2\pi} \int_0^h [\text{termD1} + \text{termX1} + \text{termP1}] r \psi_\alpha dr d\theta}_{\text{LHS}}
\tag{48}$$

1. the first  $\mathcal{D}1$  tems:

We ref the wiki [https://en.wikipedia.org/wiki/Leibniz\\_integral\\_rule](https://en.wikipedia.org/wiki/Leibniz_integral_rule)

General form: Differentiation under the integral sign:

$$\frac{d}{dx} \left( \int_{a(x)}^{b(x)} f(x, t) dt \right) - f(x, b(x)) \cdot \frac{d}{dx} b(x) + f(x, a(x)) \cdot \frac{d}{dx} a(x) = \int_{a(x)}^{b(x)} \frac{d}{dx} f(x, t) dt \tag{49}$$

$$df = \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy \tag{50}$$

For partial difference, for a given  $\beta$ , the derivation of the function  $g(\alpha) = \int_{a(\alpha)}^{b(\beta)} f(x, \alpha) dx$  is

$$\frac{d}{d\alpha} \left( \int_{a(\alpha)}^{b(\beta)} f(x, \alpha) dx \right) - 0 + \frac{da(\alpha)}{d\alpha} f(a(\alpha), \alpha) = \int_{a(\alpha)}^{b(\beta)} \frac{\partial}{\partial \alpha} f(x, \alpha) dx \tag{51}$$

1.1

$$\begin{aligned}
& \sum_{b=-\infty}^{\infty} \int_0^{2\pi} \int_0^h [-rU^{a-b} \frac{\partial U^b}{\partial s}] \psi_\alpha dr d\theta \\
&= \sum_{b=-\infty}^{\infty} - \int_0^{2\pi} \int_0^h \frac{\partial}{\partial s} [rU^{a-b} U^b \psi_\alpha] dr d\theta \\
&\quad + \int_0^{2\pi} \int_0^h U^b \frac{\partial}{\partial s} [rU^{a-b} \psi_\alpha] dr d\theta \\
&= \sum_{b=-\infty}^{\infty} - \frac{\partial}{\partial s} \int_0^{2\pi} \int_0^h [rU^{a-b} U^b \psi_\alpha] dr d\theta + \int_0^{2\pi} \frac{dh(s)}{ds} [rU^{a-b} U^b \psi_\alpha]_{r=h} d\theta \\
&\quad + \int_0^{2\pi} \int_0^h \frac{r \partial U^{a-b}}{\partial s} U^b \psi_\alpha dr d\theta \\
&\quad + \int_0^{2\pi} \int_0^h \frac{\partial \psi_\alpha}{\partial s} r U^{a-b} U^b dr d\theta
\end{aligned} \tag{52}$$

here, we gives a relationship between  $U^a$  and  $V^a$  at the boundary which to dliminate  $V^a$  tems:

$$h'U^{a-b} = (1 - \kappa h \cos \phi) V^{a-b} \tag{53}$$

1.2

$$\begin{aligned}
& \sum_{b=-\infty}^{\infty} \int_0^{2\pi} \int_0^h [-(1 - \kappa r \cos \phi) V^{a-b} \frac{\partial U^b}{\partial r}] r \psi_\alpha dr d\theta \\
&= - \sum_{b=-\infty}^{\infty} \int_0^{2\pi} \int_0^h \frac{\partial (r(1 - \kappa r \cos \phi) V^{a-b} U^b \psi_\alpha)}{\partial r} dr d\theta \\
&\quad + \int_0^{2\pi} \int_0^h U^b \frac{\partial (r(1 - \kappa r \cos \phi) V^{a-b} \psi_\alpha)}{\partial r} dr d\theta \\
&= - \sum_{b=-\infty}^{\infty} \int_0^{2\pi} [(r(1 - \kappa r \cos \phi) V^{a-b} U^b) \psi_\alpha]_0^h d\theta \\
&\quad + \int_0^{2\pi} \int_0^h U^b \frac{\partial (r(1 - \kappa r \cos \phi) V^{a-b})}{\partial r} \psi_\alpha dr d\theta \\
&\quad + \int_0^{2\pi} \int_0^h U^b \frac{\partial (\psi_\alpha)}{\partial r} r(1 - \kappa r \cos \phi) V^{a-b} dr d\theta \\
&= - \sum_{b=-\infty}^{\infty} \int_0^{2\pi} [(h h' U^{a-b} U^b) \psi_\alpha]_0^h d\theta \\
&\quad + \int_0^{2\pi} \int_0^h U^b \frac{\partial (r(1 - \kappa r \cos \phi) V^{a-b})}{\partial r} \psi_\alpha dr d\theta \\
&\quad + \int_0^{2\pi} \int_0^h U^b \frac{\partial (\psi_\alpha)}{\partial r} r(1 - \kappa r \cos \phi) V^{a-b} dr d\theta
\end{aligned} \tag{54}$$

1.3

$$\begin{aligned}
& \sum_{b=-\infty}^{\infty} \int_0^{2\pi} \int_0^h [-(1 - \kappa r \cos \phi) \frac{W^{a-b}}{r} \frac{\partial U^b}{\partial \theta}] r \psi_\alpha dr d\theta \\
&= - \sum_{b=-\infty}^{\infty} \int_0^{2\pi} \int_0^h \frac{\partial((1 - \kappa r \cos \phi) W^{a-b} U^b \psi_\alpha)}{\partial \theta} dr d\theta \\
&\quad + \int_0^{2\pi} \int_0^h U^b \frac{\partial((1 - \kappa r \cos \phi) W^{a-b})}{\partial \theta} \psi_\alpha dr d\theta \\
&\quad + \int_0^{2\pi} \int_0^h U^b \frac{\partial(\psi_\alpha)}{\partial \theta} ((1 - \kappa r \cos \phi) W^{a-b}) dr d\theta \\
&= - \sum_{b=-\infty}^{\infty} \int_0^h [(1 - \kappa r \cos \phi) W^{a-b} U^b \psi_\alpha]_0^{2\pi} dr \\
&\quad + \int_0^{2\pi} \int_0^h U^b \frac{\partial((1 - \kappa r \cos \phi) W^{a-b})}{\partial \theta} \psi_\alpha dr d\theta \\
&\quad + \int_0^{2\pi} \int_0^h U^b \frac{\partial(\psi_\alpha)}{\partial \theta} ((1 - \kappa r \cos \phi) W^{a-b}) dr d\theta \\
&= 0 \text{(periodic)} + \int_0^{2\pi} \int_0^h U^b \frac{\partial((1 - \kappa r \cos \phi) W^{a-b})}{\partial \theta} \psi_\alpha dr d\theta \\
&\quad + \int_0^{2\pi} \int_0^h U^b \frac{\partial(\psi_\alpha)}{\partial \theta} ((1 - \kappa r \cos \phi) W^{a-b}) dr d\theta
\end{aligned} \tag{55}$$

Combine together:

$$\begin{aligned}
& \frac{\int_0^{2\pi} \int_0^h [\text{term D1}] r \psi_\alpha dr d\theta}{\{\left( \int_0^{2\pi} \int_0^h \frac{\partial \psi_\alpha}{\partial s} r U^{a-b} U^b dr d\theta \right.} \\
& \quad \left. + \int_0^{2\pi} \int_0^h \frac{\partial(\psi_\alpha)}{\partial r} r (1 - \kappa r \cos \phi) V^{a-b} U^b dr d\theta \right. \\
& \quad \left. + \int_0^{2\pi} \int_0^h \frac{\partial(\psi_\alpha)}{\partial \theta} ((1 - \kappa r \cos \phi) W^{a-b} U^b) dr d\theta \right. \\
& \quad \left. + \left( \int_0^{2\pi} \int_0^h U^b \frac{\partial U^{a-b}}{\partial s} r \psi_\alpha dr d\theta \right. \right. \\
& \quad \left. \left. + \int_0^{2\pi} \int_0^h U^b \frac{\partial(r(1 - \kappa r \cos \phi) V^{a-b})}{\partial r} \psi_\alpha dr d\theta \right. \right. \\
& \quad \left. \left. + \int_0^{2\pi} \int_0^h U^b \frac{\partial((1 - \kappa r \cos \phi) W^{a-b})}{\partial \theta} \psi_\alpha dr d\theta \right) \right. \\
& \quad \left. - \frac{\partial}{\partial s} \int_0^{2\pi} \int_0^h [r U^{a-b} U^b \psi_\alpha] dr d\theta \right\}
\end{aligned} \tag{56}$$

We apply eq 5, find that:

$$\begin{aligned}
& -i(a-b)\kappa r(1 - \kappa r \cos(\phi)) P^{a-b} + \left[ \frac{\partial(U^{a-b}r)}{\partial s} + \frac{\partial(V^{a-b}r(1 - \kappa r \cos(\phi)))}{\partial r} + \frac{\partial(W^{a-b}(1 - \kappa r \cos(\phi)))}{\partial \theta} \right] \\
&= o(M^2)
\end{aligned} \tag{57}$$

We have the second terms in eq(31) are equal to:

$$\begin{aligned} & \int_0^{2\pi} \int_0^h U^b [-i(a-b)\kappa r(1-\kappa r \cos(\phi)) P^{a-b}] \psi_\alpha dr d\theta \\ &= i(a-b)\kappa \Psi_{\alpha\beta\gamma} [r(1-\kappa r \cos(\phi))] P_\beta^{a-b} U_\gamma^b \end{aligned} \quad (58)$$

And, the longitudinal derivation  $s$  can also be expand about the duct modes, with note

$[r], (\theta), \{s\}$ :

$$\begin{aligned} & \frac{\partial}{\partial s} \int_0^{2\pi} \int_0^h [r U^{a-b} U^b \psi_\alpha] dr d\theta \\ &= \frac{\partial}{\partial s} \int_0^{2\pi} \int_0^h [r \psi_\beta \psi_\gamma \psi_\alpha] U_\beta^{a-b} U_\gamma^b dr d\theta \\ &= \frac{\partial}{\partial s} \left( \int_0^{2\pi} \int_0^h [r \psi_\beta \psi_\gamma \psi_\alpha] dr d\theta \right) U_\beta^{a-b} U_\gamma^b \\ &+ \frac{\partial U_\beta^{a-b} U_\gamma^b}{\partial s} \int_0^{2\pi} \int_0^h [r \psi_\beta \psi_\gamma \psi_\alpha] dr d\theta \\ &= \frac{\partial}{\partial s} \left( \int_0^{2\pi} \int_0^h [r \psi_\beta \psi_\gamma \psi_\alpha] dr d\theta \right) U_\beta^{a-b} U_\gamma^b \\ &+ \left( \frac{dU_\beta^{a-b}}{ds} U_\gamma^b + \frac{dU_\gamma^b}{ds} U_\beta^{a-b} \right) \int_0^{2\pi} \int_0^h [r \psi_\beta \psi_\gamma \psi_\alpha] dr d\theta \end{aligned} \quad (59)$$

2. the second  $\mathcal{X}1$  tems:

$$\begin{aligned} term \mathcal{X}1 : & \sum_{b=-\infty}^{\infty} \kappa \cos(\phi) U^{a-b} V^b - \kappa \sin(\phi) U^{a-b} W^b \\ & \sum_{b=-\infty}^{\infty} \int_0^{2\pi} \int_0^h [\kappa \cos(\phi) U^{a-b} V^b - \kappa \sin(\phi) U^{a-b} W^b] r \psi_\alpha dr d\theta \\ &= \kappa \Psi_{\alpha\beta\gamma} [\cos(\phi) U_\beta^{a-b} V_\gamma^b - \kappa \sin(\phi) U_\beta^{a-b} W_\gamma^b] \end{aligned} \quad (60)$$

3. the second  $\mathcal{P}1$  tems:

$$\begin{aligned} term \mathcal{P}1 : & \sum_{b=-\infty}^{\infty} i b \kappa (1 - \kappa r \cos(\phi)) P^{a-b} U^b \\ & \sum_{b=-\infty}^{\infty} \int_0^{2\pi} \int_0^h [i b \kappa (1 - \kappa r \cos(\phi)) P^{a-b} U^b] r \psi_\alpha dr d\theta \\ &= i b \kappa \Psi_{\alpha\beta\gamma} [r (1 - \kappa r \cos(\phi))] P_\beta^{a-b} U_\gamma^b \end{aligned} \quad (61)$$

4. The LHS terms:

$$\frac{\partial P^a}{\partial s} - i a \kappa (1 - \kappa r \cos(\phi)) U^a$$

From 1.1 as example, we know that

$$\begin{aligned}
& \int_0^{2\pi} \int_0^h \frac{\partial}{\partial s} [rU^{a-b}U^b\psi_\alpha] dr d\theta \\
&= \frac{\partial}{\partial s} \int_0^{2\pi} \int_0^h [rU^{a-b}U^b\psi_\alpha] dr d\theta - \int_0^{2\pi} \frac{dh(s)}{ds} [rU^{a-b}U^b\psi_\alpha]_{r=h} d\theta
\end{aligned} \tag{62}$$

4.1

$$\begin{aligned}
& \int_0^{2\pi} \int_0^h \left[ \frac{\partial P^a}{\partial s} \right] r\psi_\alpha dr d\theta \\
&= \int_0^{2\pi} \int_0^h \left[ \frac{\partial (P_\beta^a \psi_\beta)}{\partial s} \right] r\psi_\alpha dr d\theta \\
&= \int_0^{2\pi} \int_0^h \left[ \frac{\partial (P_\beta^a \psi_\beta)}{\partial s} r\psi_\alpha \right] dr d\theta - \int_0^{2\pi} \int_0^h \left[ \frac{\partial (\psi_\alpha)}{\partial s} \right] r\psi_\beta dr d\theta P_\beta^a \\
&= \frac{\partial}{\partial s} \int_0^{2\pi} \int_0^h (P_\beta^a \psi_\beta) r\psi_\alpha dr d\theta - \frac{dh(s)}{ds} [P_\beta^a \psi_\beta r\psi_\alpha]_{r=h} - \int_0^{2\pi} \int_0^h \left[ \frac{\partial (\psi_\alpha)}{\partial s} \right] r\psi_\beta dr d\theta P_\beta^a \\
&= \frac{d}{ds} (P_\beta^a \hat{\delta}_{\alpha\beta}) - \int_0^{2\pi} \frac{dh(s)}{ds} [P_\beta^a \psi_\beta r\psi_\alpha]_{r=h} d\theta - \int_0^{2\pi} \int_0^h \left[ \frac{\partial (\psi_\alpha)}{\partial s} \right] r\psi_\beta dr d\theta P_\beta^a
\end{aligned} \tag{63}$$

4.2

$$\begin{aligned}
& \int_0^{2\pi} \int_0^h [-ia\kappa(1 - \kappa r \cos\phi) U^a] r\psi_\alpha dr d\theta \\
&= -ia\kappa \Psi_{\alpha\beta} [r(1 - \kappa r \cos\phi)] U_\beta^a
\end{aligned} \tag{64}$$

Finally, putting all together becomes:

$$\begin{aligned}
& \frac{d}{ds}(P_\beta^a)\widehat{\delta}_{\alpha\beta} - \int_0^{2\pi} \frac{dh(s)}{ds} [P_\beta^a \psi_\beta r \psi_\alpha]_{r=h} d\theta - \int_0^{2\pi} \int_0^h [\frac{\partial(\psi_\alpha)}{\partial s}] r \psi_\beta dr d\theta P_\beta^a - ia\kappa\Psi_{\alpha\beta}[r(1-\kappa r\cos\phi)]U_\beta^a \\
&= \frac{d}{ds}(P_\beta^a)\widehat{\delta}_{\alpha\beta} - ia\kappa\Psi_{\alpha\beta}[r(1-\kappa r\cos\phi)]U_\beta^a - \int_0^{2\pi} h h' [P_\beta^a \psi_\beta \psi_\alpha]_{r=h} d\theta - \Psi_{\{\alpha\}\beta}[r]P_\beta^a \\
&= \sum_{b=-\infty}^{\infty} \\
&\quad (eq31) : \{ (\int_0^{2\pi} \int_0^h \frac{\partial\psi_\alpha}{\partial s} r U^{a-b} U^b dr d\theta \\
&\quad + \int_0^{2\pi} \int_0^h \frac{\partial(\psi_\alpha)}{\partial r} r(1-\kappa r\cos\phi) V^{a-b} U^b dr d\theta \\
&\quad + (eq33) : i(a-b)\kappa\Psi_{\alpha\beta\gamma}[r(1-\kappa r\cos\phi)]P_\beta^{a-b} U_\gamma^b \\
&\quad eq(34) : -\frac{\partial}{\partial s} (\int_0^{2\pi} \int_0^h [r \psi_\beta \psi_\gamma \psi_\alpha] dr d\theta) U_\beta^{a-b} U_\gamma^b \\
&\quad - (\frac{dU_\beta^{a-b}}{ds} U_\gamma^b + \frac{dU_\gamma^b}{ds} U_\beta^{a-b}) \int_0^{2\pi} \int_0^h [r \psi_\beta \psi_\gamma \psi_\alpha] dr d\theta \\
&\quad + eq(35) : \kappa\Psi_{\alpha\beta\gamma}[r\cos\phi] U_\beta^{a-b} V_\gamma^b - \kappa\Psi_{\alpha\beta\gamma}[r\sin\phi] U_\beta^{a-b} W_\gamma^b \\
&\quad + eq(36) : ib\kappa\Psi_{\alpha\beta\gamma}[r(1-\kappa r\cos\phi)]P_\beta^{a-b} U_\gamma^b \\
&\quad = (abbreviation) : \\
&\quad \Psi_{\{\alpha\}\beta\gamma}[r] U_\beta^{a-b} U_\gamma^a + \Psi_{[\alpha]\beta\gamma}[r(1-\kappa r\cos\phi)] V_\beta^{a-b} U_\gamma^a + \Psi_{(\alpha)\beta\gamma}[(1-\kappa r\cos\phi)] W_\beta^{a-b} U_\gamma^a \\
&\quad + i(a-b)\kappa\Psi_{\alpha\beta\gamma}[r(1-\kappa r\cos\phi)]P_\beta^{a-b} U_\gamma^b \\
&\quad - \frac{\partial}{\partial s} (\Psi_{\alpha\beta\gamma}[r] U_\beta^{a-b} U_\gamma^b - \Psi_{\alpha\beta\gamma}[r] (\frac{dU_\beta^{a-b}}{ds} U_\gamma^b + \frac{dU_\gamma^b}{ds} U_\beta^{a-b})) \\
&\quad + \kappa\Psi_{\alpha\beta\gamma}[r\cos\phi] U_\beta^{a-b} V_\gamma^b - \kappa\Psi_{\alpha\beta\gamma}[r\sin\phi] U_\beta^{a-b} W_\gamma^b \\
&\quad + ib\kappa\Psi_{\alpha\beta\gamma}[r(1-\kappa r\cos\phi)]P_\beta^{a-b} U_\gamma^b
\end{aligned} \tag{65}$$

with the  $e^s$  term:

$$\begin{aligned}
& -ia\kappa U^a + \frac{1}{1-\kappa r\cos\phi} \frac{\partial P^a}{\partial s} \\
&= \sum_{b=-\infty}^{\infty} term\mathcal{D}1 : -\frac{U^{a-b}}{1-\kappa r\cos\phi} \frac{\partial U^b}{\partial s} - V^{a-b} \frac{\partial U^b}{\partial r} - \frac{W^{a-b}}{r} \frac{\partial U^b}{\partial \theta} \\
&+ \sum_{b=-\infty}^{\infty} term\mathcal{X}1 : \frac{\kappa\cos\phi}{1-\kappa r\cos\phi} U^{a-b} V^b - \frac{\kappa\sin\phi}{(1-\kappa r\cos\phi)} U^{a-b} W^b \\
&+ \sum_{b=-\infty}^{\infty} term\mathcal{P}1 : ib\kappa P^{a-b} U^b
\end{aligned} \tag{66}$$

We have:

$$\begin{aligned}
& -ia\kappa\Psi_{\alpha\beta}[r(1-\kappa r\cos\phi)]U_\beta^a + \frac{d}{ds}(P_\beta^a)\widehat{\delta}_{\alpha\beta} - \int_0^{2\pi} hh'[\psi_\beta\psi_\alpha]_{r=h}d\theta P_\beta^a - \Psi_{\{\alpha\}\beta}[r]P_\beta^a \\
& = \underline{\text{termD1}} : \Psi_{\{\alpha\}\beta\gamma}[r]U_\beta^{a-b}U_\gamma^a + \Psi_{[\alpha]\beta\gamma}[r(1-\kappa r\cos\phi)]V_\beta^{a-b}U_\gamma^a + \Psi_{(\alpha)\beta\gamma}[(1-\kappa r\cos\phi)]W_\beta^{a-b}U_\gamma^a \\
& \quad + \underline{\text{term}(D1+\mathcal{P}1)} : i(a)\kappa\Psi_{\alpha\beta\gamma}[r(1-\kappa r\cos\phi)]P_\beta^{a-b}U_\gamma^b \\
& \quad - \underline{\text{termD1}} : \frac{\partial}{\partial s}(\Psi_{\alpha\beta\gamma}[r]U_\beta^{a-b}U_\gamma^b - \Psi_{\alpha\beta\gamma}[r](\frac{dU_\beta^{a-b}}{ds}U_\gamma^b + \frac{dU_\gamma^b}{ds}U_\beta^{a-b})) \\
& \quad + \underline{\text{termX1}} : \kappa\Psi_{\alpha\beta\gamma}[r\cos\phi]U_\beta^{a-b}V_\gamma^b - \kappa\Psi_{\alpha\beta\gamma}[r\sin\phi]U_\beta^{a-b}W_\gamma^b
\end{aligned} \tag{67}$$

## B. Momentum $e^r$ term

Second, deal with the  $e^r$  term:

$$\begin{aligned}
& -ia\kappa V^a + \frac{\partial P^a}{\partial r} \\
& = \sum_{b=-\infty}^{\infty} \text{termD2} : -\frac{U^{a-b}}{1-\kappa r\cos\phi} \frac{\partial V^b}{\partial s} - V^{a-b} \frac{\partial V^b}{\partial r} - \frac{W^{a-b}}{r} \frac{\partial V^b}{\partial \theta} \\
& \quad + \sum_{b=-\infty}^{\infty} \text{termX2} : \frac{W^{a-b}W^b}{r} - \frac{\kappa\cos\phi}{(1-\kappa r\cos\phi)} U^{a-b}U^b \\
& \quad + \text{termP2} : \sum_{b=-\infty}^{\infty} ib\kappa P^{a-b}V^b
\end{aligned} \tag{68}$$

$$\text{LHS-2: } \frac{\partial P^a}{\partial r}(1-\kappa r\cos\phi)$$

$$\begin{aligned}
& \int_0^{2\pi} \int_0^h [\frac{\partial P^a}{\partial r}(1-\kappa r\cos\phi)]r\psi_\alpha dr d\theta \\
& = \int_0^{2\pi} \int_0^h [\frac{\partial \psi_\beta(1-\kappa r\cos\phi)r\psi_\alpha}{\partial r}]dr d\theta P_\beta^a - \int_0^{2\pi} \int_0^h [\frac{\partial \psi_\alpha(1-\kappa r\cos\phi)r}{\partial r}]\psi_\beta dr d\theta P_\beta^a \\
& \quad = \int_0^{2\pi} \int_0^h [\frac{\partial \psi_\beta(1-\kappa r\cos\phi)r\psi_\alpha}{\partial r}]dr d\theta P_\beta^a \\
& \quad - \int_0^{2\pi} \int_0^h [\frac{\partial \psi_\alpha}{\partial r}](1-\kappa r\cos\phi)r\psi_\beta dr d\theta P_\beta^a - \int_0^{2\pi} \int_0^h [\frac{\partial (1-\kappa r\cos\phi)r}{\partial r}]\psi_\alpha\psi_\beta dr d\theta P_\beta^a \\
& \quad = \int_0^{2\pi} [\psi_\alpha\psi_\beta r(1-\kappa r\cos\phi)]_0^h d\theta P_\beta^a \\
& \quad - \int_0^{2\pi} \int_0^h [\frac{\partial \psi_\alpha}{\partial r}](1-\kappa r\cos\phi)r\psi_\beta dr d\theta P_\beta^a - \int_0^{2\pi} \int_0^h [1-2\kappa r\cos\phi]\psi_\alpha\psi_\beta dr d\theta P_\beta^a \\
& = \int_0^{2\pi} [\psi_\alpha\psi_\beta r(1-\kappa r\cos\phi)]_0^h d\theta P_\beta^a - \Psi_{[\alpha]\beta}[r(1-\kappa r\cos\phi)]P_\beta^a - \Psi_{\alpha\beta}[1-2\kappa r\cos\phi]P_\beta^a
\end{aligned} \tag{69}$$

The derivation of  $\text{termD2}$  is identical to A, we are not prove it again.  $\text{TermP2}$  also could be combine with the part separated term of  $\text{termD2}$  with  $V^b$ .  $\text{TermX2}$  is also easy to derive.

Thus, we have the final equation:

$$\begin{aligned}
& -ia\kappa\Psi_{\alpha\beta}[r(1-\kappa r\cos\phi)]V_\beta^a \\
& \int_0^{2\pi} [\psi_\alpha\psi_\beta r(1-\kappa r\cos\phi)]_0^h d\theta P_\beta^a - \Psi_{[\alpha]\beta}[r(1-\kappa r\cos\phi)]P_\beta^a - \Psi_{\alpha\beta}[1-2\kappa r\cos\phi]P_\beta^a \\
& = \underline{\text{termD2}} : \Psi_{\{\alpha\}\beta\gamma}[r]U_\beta^{a-b}V_\gamma^a + \Psi_{[\alpha]\beta\gamma}[r(1-\kappa r\cos\phi)]V_\beta^{a-b}V_\gamma^a + \Psi_{(\alpha)\beta\gamma}[(1-\kappa r\cos\phi)]W_\beta^{a-b}V_\gamma^a \\
& \quad + \underline{\text{term}(D2 + \mathcal{P}2)} : i(a)\kappa\Psi_{\alpha\beta\gamma}[r(1-\kappa r\cos\phi)]P_\beta^{a-b}V_\gamma^b \\
& \quad + \underline{\text{termD2}} : -\frac{\partial}{\partial s}(\Psi_{\alpha\beta\gamma}[r]U_\beta^{a-b}V_\gamma^b - \Psi_{\alpha\beta\gamma}[r](\frac{dU_\beta^{a-b}}{ds}V_\gamma^b + \frac{dV_\gamma^b}{ds}U_\beta^{a-b})) \\
& \quad + \underline{\text{termX2}} : \Psi_{\alpha\beta\gamma}[1-\kappa r\cos\phi]W_\beta^{a-b}W_\gamma^b - \kappa\Psi_{\alpha\beta\gamma}[r\cos\phi]U_\beta^{a-b}U_\gamma^b
\end{aligned} \tag{70}$$

### C. Momentum $e^\theta$ term

Third, deal with the  $e^\theta$  term:

$$\begin{aligned}
& -ia\kappa W^a + \frac{1}{r}\frac{\partial P^a}{\partial\theta} \\
& = \sum_{b=-\infty}^{\infty} \text{termD3} : -\frac{U^{a-b}}{1-\kappa r\cos\phi}\frac{\partial W^b}{\partial s} - V^{a-b}\frac{\partial W^b}{\partial r} - \frac{W^{a-b}}{r}\frac{\partial W^b}{\partial\theta} \\
& \quad + \sum_{b=-\infty}^{\infty} \text{termX3} : \frac{\kappa\sin\phi}{(1-\kappa r\cos\phi)}U^{a-b}U^b - \frac{W^{a-b}V^b}{r} \\
& \quad + \text{termP3} : \sum_{b=-\infty}^{\infty} ib\kappa P^{a-b}W^b
\end{aligned} \tag{71}$$

$$\begin{aligned}
& \text{LHS-2: } \frac{\partial P^a}{\partial\theta} \frac{(1-\kappa r\cos\phi)}{r} \\
& \quad \int_0^{2\pi} \int_0^h [\frac{\partial P^a}{\partial\theta} \frac{(1-\kappa r\cos\phi)}{r}] r\psi_\alpha dr d\theta \\
& = \int_0^{2\pi} \int_0^h [\frac{\partial\psi_\beta(1-\kappa r\cos\phi)\psi_\alpha}{\partial\theta}] dr d\theta P_\beta^a - \int_0^{2\pi} \int_0^h [\frac{\partial\psi_\alpha(1-\kappa r\cos\phi)}{\partial\theta}] \psi_\beta dr d\theta P_\beta^a \\
& = 0 - \int_0^{2\pi} \int_0^h [\frac{\partial\psi_\alpha}{\partial\theta}](1-\kappa r\cos\phi)\psi_\beta dr d\theta P_\beta^a - \int_0^{2\pi} \int_0^h [\frac{\partial(1-\kappa r\cos\phi)}{\partial\theta}] \psi_\alpha\psi_\beta dr d\theta P_\beta^a \\
& = - \int_0^{2\pi} \int_0^h [\frac{\partial\psi_\alpha}{\partial\theta}](1-\kappa r\cos\phi)\psi_\beta dr d\theta P_\beta^a + \kappa \int_0^{2\pi} \int_0^h [r\sin\phi]\psi_\alpha\psi_\beta dr d\theta P_\beta^a \\
& = -\Psi_{(\alpha)\beta}[(1-\kappa r\cos\phi)]P_\beta^a - \kappa\Psi_{\alpha\beta}[r\sin\phi]P_\beta^a
\end{aligned} \tag{72}$$

The derivation of  $\text{termD3}$  is identical to A, we are not prove it again.  $\text{TermP3}$  also could be combine with the part separated term of  $\text{termD3}$  with  $W^b$ .  $\text{TermX3}$  is also easy to derive.

Thus, we have the final equation:

$$\begin{aligned}
& -ia\kappa\Psi_{\alpha\beta}[r(1-\kappa r\cos\phi)]W_\beta^a \\
& -\Psi_{(\alpha)\beta}[(1-\kappa r\cos\phi)]P_\beta^a - \kappa\Psi_{\alpha\beta}[r\sin\phi]P_\beta^a \\
= & \underline{\text{termD2}} : \Psi_{\{\alpha\}\beta\gamma}[r]U_\beta^{a-b}W_\gamma^a + \Psi_{[\alpha]\beta\gamma}[r(1-\kappa r\cos\phi)]V_\beta^{a-b}W_\gamma^a + \Psi_{(\alpha)\beta\gamma}[(1-\kappa r\cos\phi)]W_\beta^{a-b}W_\gamma^a \\
& + \underline{\text{term}(D2+P2)} : i(a)\kappa\Psi_{\alpha\beta\gamma}[r(1-\kappa r\cos\phi)]P_\beta^{a-b}W_\gamma^b \\
& + \underline{\text{termD2}} : -\frac{\partial}{\partial s}(\Psi_{\alpha\beta\gamma}[r]U_\beta^{a-b}V_\gamma^b - \Psi_{\alpha\beta\gamma}[r](\frac{dU_\beta^{a-b}}{ds}W_\gamma^b + \frac{dW_\gamma^b}{ds}U_\beta^{a-b})) \\
& + \underline{\text{termX2}} : \kappa\Psi_{\alpha\beta\gamma}[r\sin\phi]U_\beta^{a-b}U_\gamma^b - \Psi_{\alpha\beta\gamma}[1-\kappa r\cos\phi]W_\beta^{a-b}V_\gamma^b
\end{aligned} \tag{73}$$

#### IV. Merge the four equations and eliminate the $V_\gamma^b$ and $W_\gamma^b$

##### A. $V_\alpha^a$ & $W_\alpha^a$ for RHS

Using the linear relationships:

$$\begin{aligned}
ia\kappa V^a &= \frac{\partial P^a}{\partial r} \\
&:= \int \int ia k V_\beta^a \psi_\beta r \psi_\alpha dr d\theta = \int \int \frac{\partial P_\beta^a \psi_\beta}{\partial r} r \psi_\alpha dr d\theta \\
= ia\kappa V_\beta^a \hat{\delta}_{\alpha\beta} &= \Psi_{\alpha[\beta]}[r]P_\beta^a = \int_0^{2\pi} [r\psi_\alpha \psi_\beta]_0^h d\theta P_\beta^a - \int \int \psi_\alpha \psi_\beta dr d\theta P_\beta^a - \int \int \frac{\partial \psi_\alpha}{\partial r} \psi_\beta r dr d\theta P_\beta^a
\end{aligned} \tag{74}$$

$$\begin{aligned}
ia\kappa W^a &= \frac{1}{r} \frac{\partial P^a}{\partial \theta} \\
&:= \int \int ia k W_\beta^a \psi_\beta r \psi_\alpha dr d\theta = \int \int \frac{\partial P_\beta^a \psi_\beta}{\partial \theta} \frac{1}{r} r \psi_\alpha dr d\theta \\
= ia\kappa W_\beta^a \hat{\delta}_{\alpha\beta} &= \Psi_{\alpha(\beta)}[r]P_\beta^a = 0 - \Psi_{(\alpha)\beta}[r]P_\beta^a
\end{aligned} \tag{75}$$

Thus, we can establish relationships between the transverse modes and pressure modes (no summation over a)

$$\underline{\underline{V_\beta^a \hat{\delta}_{\alpha\beta} = \frac{1}{ia\kappa} [\int_0^{2\pi} [r\psi_\alpha \psi_\beta]_0^h d\theta - \Psi_{\alpha\beta} - \Psi_{[\alpha]\beta}[r]] P_\beta^a = \mathbf{V}_{\alpha\beta}^a P_\beta^a]}} \tag{76}$$

$$\underline{\underline{W_\beta^a \hat{\delta}_{\alpha\beta} = -\frac{1}{ia\kappa} \Psi_{(\alpha)\beta} P_\beta^a = \mathbf{W}_{\alpha\beta}^a P_\beta^a}}} \tag{77}$$

**B.**  $\frac{d}{ds}V_\alpha^a$  &  $\frac{d}{ds}W_\alpha^a$  for RHS

We also require modal expressions for  $\frac{d}{ds}V_\alpha^a$  and  $\frac{d}{ds}W_\alpha^a$ .

We differentiate eq71 with respect to s:

$$\begin{aligned}\frac{\partial V^a}{\partial s} &= \frac{1}{ia\kappa} \frac{\partial^2 P^a}{\partial s \partial r} \\ &= \frac{\partial}{\partial r}((1 - \kappa r \cos \phi) U^a)\end{aligned}\quad (78)$$

where we have used symmetry of mixed partials and the linear expression for  $\frac{\partial P^a}{\partial s}$  from eq 21.

From 1.1 as example, we know that

$$\begin{aligned}&\int_0^{2\pi} \int_0^h \frac{\partial}{\partial s} [r U^{a-b} U^b \psi_\alpha] dr d\theta \\ &= \frac{\partial}{\partial s} \int_0^{2\pi} \int_0^h [r U^{a-b} U^b \psi_\alpha] dr d\theta - \int_0^{2\pi} \frac{dh(s)}{ds} [r U^{a-b} U^b \psi_\alpha]_{r=h} d\theta\end{aligned}\quad (79)$$

here, we gives a relationship between  $U^a$  and  $V^a$  at the boundary which to dliminate  $V^a$  tems:

$$h' U_\beta^a = (1 - \kappa h \cos \phi) V_\beta^a \quad (80)$$

Multiplying this expression by  $r \phi_\alpha$  and integrating across section of the duct, we obtain:

$$\begin{aligned}&\int_0^{2\pi} \int_0^h \frac{\partial V^a}{\partial s} r \psi_\alpha dr d\theta = \int_0^{2\pi} \int_0^h \frac{\partial}{\partial r} ((1 - \kappa r \cos \phi) U^a) r \psi_\alpha dr d\theta \\ LHS &:= \int_0^{2\pi} \int_0^h \frac{\partial [V_\beta^a \psi_\beta r \psi_\alpha]}{\partial s} dr d\theta - \int_0^{2\pi} \int_0^h \frac{\partial r \psi_\alpha}{\partial s} \psi_\beta dr d\theta V_\beta^a \\ &= \frac{\partial}{\partial s} \int_0^{2\pi} \int_0^h [V_\beta^a \psi_\beta r \psi_\alpha] dr d\theta - \int_0^{2\pi} \frac{dh(s)}{ds} [V_\beta^a \psi_\beta r \psi_\alpha]_0^h d\theta - \int_0^{2\pi} \int_0^h \frac{\partial \psi_\alpha}{\partial s} \psi_\beta r dr d\theta V_\beta^a \\ &= \frac{d}{ds} V_\beta^a \hat{\delta}_{\alpha\beta} - \int_0^{2\pi} \frac{h'^2}{1 - \kappa h \cos \phi} [r \psi_\beta \psi_\alpha]_0^h d\theta U_\beta^a - \Psi_{\{\alpha\}\beta} [r] V_\beta^a \\ RHS &:= \int_0^{2\pi} \int_0^h \frac{\partial}{\partial r} ((1 - \kappa r \cos \phi) U_\beta^a \psi_\beta r \psi_\alpha) dr d\theta - \int_0^{2\pi} \int_0^h \frac{\partial (r \psi_\alpha)}{\partial r} (1 - \kappa r \cos \phi) U_\beta^a \psi_\beta dr d\theta \\ &= \int_0^{2\pi} [r (1 - \kappa r \cos \phi) \psi_\beta \psi_\alpha]_0^h d\theta U_\beta^a - \int_0^{2\pi} \int_0^h \frac{\partial \psi_\alpha}{\partial r} r (1 - \kappa r \cos \phi) \psi_\beta dr d\theta U_\beta^a - \int_0^{2\pi} \int_0^h (1 - \kappa r \cos \phi) \psi_\alpha \psi_\beta dr d\theta U_\beta^a \\ &= \int_0^{2\pi} [r (1 - \kappa r \cos \phi) \psi_\beta \psi_\alpha]_0^h d\theta U_\beta^a - \Psi_{[\alpha]\beta} [r (1 - \kappa r \cos \phi)] U_\beta^a - \Psi_{\alpha\beta} [(1 - \kappa r \cos \phi)] U_\beta^a\end{aligned}\quad (81)$$

Thus, LHS=RHS, we have:

$$\begin{aligned}\frac{d}{ds} V_\beta^a \hat{\delta}_{\alpha\beta} &= \int_0^{2\pi} \frac{h'^2}{1 - \kappa h \cos \phi} [r \psi_\beta \psi_\alpha]_0^h d\theta U_\beta^a + \Psi_{\{\alpha\}\beta} [r] V_\beta^a \\ &+ \int_0^{2\pi} [r (1 - \kappa r \cos \phi) \psi_\beta \psi_\alpha]_0^h d\theta U_\beta^a - \Psi_{[\alpha]\beta} [r (1 - \kappa r \cos \phi)] U_\beta^a - \Psi_{\alpha\beta} [(1 - \kappa r \cos \phi)] U_\beta^a\end{aligned}\quad (82)$$

Similarly for  $W^a$ , differentiating eq50 with respect to  $s$  and substituting the linear expression for  $\frac{\partial P^a}{\partial s}$  by eq21:

$$\begin{aligned}\frac{\partial W^a}{\partial s} &= \frac{1}{ia\kappa} \frac{\partial^2 P^a}{\partial s \partial \theta} \\ &= \frac{1}{r} \frac{\partial}{\partial \theta} ((1 - \kappa r \cos \phi) U^a)\end{aligned}\quad (83)$$

Multiplying this expression by  $r\phi_\alpha$  and integrating across section of the duct, we obtain:

$$\begin{aligned}\int_0^{2\pi} \int_0^h \frac{\partial W^a}{\partial s} r \psi_\alpha dr d\theta &= \int_0^{2\pi} \int_0^h \frac{1}{r} \frac{\partial}{\partial \theta} ((1 - \kappa r \cos \phi) U^a) r \psi_\alpha dr d\theta \\ LHS &:= \int_0^{2\pi} \int_0^h \frac{\partial [W_\beta^a \psi_\beta r \psi_\alpha]}{\partial s} dr d\theta - \int_0^{2\pi} \int_0^h \frac{\partial r \psi_\alpha}{\partial s} \psi_\beta dr d\theta W_\beta^a \\ &= \frac{\partial}{\partial s} \int_0^{2\pi} \int_0^h [W_\beta^a \psi_\beta r \psi_\alpha] dr d\theta - \int_0^{2\pi} \frac{dh(s)}{ds} [W_\beta^a \psi_\beta r \psi_\alpha]_0^h d\theta - \int_0^{2\pi} \int_0^h \frac{\partial \psi_\alpha}{\partial s} \psi_\beta r dr d\theta W_\beta^a \\ &= \frac{d}{ds} W_\beta^a \hat{\delta}_{\alpha\beta} - \int_0^{2\pi} \frac{dh(s)}{ds} [\psi_\beta r \psi_\alpha]_0^h d\theta W_\beta^a - \Psi_{\{\alpha\}\beta}[r] W_\beta^a \\ RHS &:= \int_0^{2\pi} \int_0^h \frac{\partial}{\partial \theta} ((1 - \kappa r \cos \phi) U_\beta^a \psi_\beta \psi_\alpha) dr d\theta - \int_0^{2\pi} \int_0^h \frac{\partial (\psi_\alpha)}{\partial \theta} (1 - \kappa r \cos \phi) U_\beta^a \psi_\beta dr d\theta \\ &= 0(\text{periodic}) - \Psi_{(\alpha)\beta}[1 - \kappa r \cos \phi] U_\beta^a\end{aligned}\quad (84)$$

Thus, LHS=RHS, we have:

$$\frac{d}{ds} W_\beta^a \hat{\delta}_{\alpha\beta} = \int_0^{2\pi} \frac{dh(s)}{ds} [\psi_\beta r \psi_\alpha]_0^h d\theta W_\beta^a + \Psi_{\{\alpha\}\beta}[r] W_\beta^a - \Psi_{(\alpha)\beta}[1 - \kappa r \cos \phi] U_\beta^a = -\Psi_{\alpha\{\beta\}}[r] W_\beta^a - \Psi_{(\alpha)\beta}[1 - \kappa r \cos \phi] U_\beta^a\quad (85)$$

## V. Substitue pressure modes for transverse velocity modes

### A. mass equation

$$\begin{aligned}\frac{dU_\beta^a}{ds} \hat{\delta}_{\alpha\beta} - \Psi_{\{\alpha\}\beta}[r] U_\beta^a - ia\kappa \Psi_{\alpha\beta}[r(1 - \kappa \cos(\phi))] P_\beta^a - \Psi_{[\alpha]\beta}[r(1 - \kappa r \cos(\phi))] \underline{\underline{V_\beta^a}} - \Psi_{(\alpha)\beta}[1 - \kappa r \cos(\phi)] \underline{\underline{W_\beta^a}} \\ = \Psi_{\alpha\beta\gamma}[r(1 - \kappa r \cos(\phi))] \sum_{b=-\infty}^{+\infty} (-ib\kappa P_\beta^{a-b} P_\gamma^b - ib\kappa U_\beta^{a-b} U_\gamma^b - ib\kappa \underline{\underline{V_\beta^{a-b}}} \underline{\underline{V_\gamma^b}} - ib\kappa \underline{\underline{W_\beta^{a-b}}} \underline{\underline{W_\gamma^b}} - ia\kappa \frac{B}{2A} P_\beta^{a-b} P_\gamma^b)\end{aligned}\quad (86)$$

Transform:

$$\frac{dU_\beta^a}{ds} \widehat{\delta}_{\alpha\beta} := \widehat{\mathbf{I}_{\alpha\beta}^a} u'_\beta$$

$$-\Psi_{\{\alpha\}\beta}[r]U_\beta^a := -\underline{\Psi_{\{\alpha\}\beta}[r]u_\beta^a} \rightarrow \mathcal{G}$$

$$\begin{cases} -ia\kappa\Psi_{\alpha\beta}[r(1-\kappa\cos(\phi))]P_\beta^a := \sum_{\beta=0}^{+\infty} -ia\kappa\Psi_{\alpha\beta}[r(1-\kappa\cos(\phi))]p_\beta^a \rightarrow \mathcal{M}_1 \\ -\Psi_{[\alpha]\beta}[r(1-\kappa r\cos(\phi))]V_\beta^a := \sum_{\beta=0}^{+\infty} -\Psi_{[\alpha]\delta}[r(1-\kappa r\cos(\phi))]\mathbf{V}_{\delta\beta}^a p_\beta^a \rightarrow \mathcal{M}_2 + \Psi_{[\alpha]\delta}[r(1-krcos\phi)](N^{-1})(o(M_2^2)) \\ -\Psi_{(\alpha)\beta}[(1-\kappa r\cos(\phi))]W_\beta^a := \sum_{\beta=0}^{+\infty} -\Psi_{(\alpha)\delta}[(1-\kappa r\cos(\phi))]\mathbf{W}_{\delta\beta}^a p_\beta^a \rightarrow \mathcal{M}_3 + \Psi_{[\alpha]\delta}[r(1-krcos\phi)](N^{-1})(o(M_3^2)) \\ \Psi_{\alpha\beta\gamma}[r(1-\kappa r\cos(\phi))] \sum_{b=-\infty}^{+\infty} (-ib\kappa P_\beta^{a-b} P_\gamma^b) := \sum_{b=-\infty}^{+\infty} -ib\kappa \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \underline{\Psi_{\alpha\beta\gamma}[(1-\kappa r\cos(\phi))]p_\beta^{a-b} p_\gamma^b} \rightarrow \mathcal{B}_2 \\ \Psi_{\alpha\beta\gamma}[r(1-\kappa r\cos(\phi))] \sum_{b=-\infty}^{+\infty} (-ib\kappa U_\beta^{a-b} U_\gamma^b) := \sum_{b=-\infty}^{+\infty} -ib\kappa \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \underline{\Psi_{\alpha\beta\gamma}[(1-\kappa r\cos(\phi))]u_\beta^{a-b} u_\gamma^b} \rightarrow \mathcal{A}_1 \\ \Psi_{\alpha\beta\gamma}[r(1-\kappa r\cos(\phi))] \sum_{b=-\infty}^{+\infty} (-ib\kappa V_\beta^{a-b} V_\gamma^b) := \sum_{b=-\infty}^{+\infty} -ib\kappa \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \sum_{\delta,\epsilon=0}^{\infty} \underline{\Psi_{\alpha\delta\epsilon}[r(1-\kappa r\cos\phi)]\mathbf{V}_{\delta\beta}^{a-b} \mathbf{V}_{\epsilon\gamma}^b p_\beta^{a-b} p_\gamma^b} \rightarrow \mathcal{B}_3 \\ \Psi_{\alpha\beta\gamma}[r(1-\kappa r\cos(\phi))] \sum_{b=-\infty}^{+\infty} (-ib\kappa W_\beta^{a-b} W_\gamma^b) := \sum_{b=-\infty}^{+\infty} -ib\kappa \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \sum_{\delta,\epsilon=0}^{\infty} \underline{\Psi_{\alpha\delta\epsilon}[r(1-\kappa r\cos\phi)]\mathbf{W}_{\delta\beta}^{a-b} \mathbf{W}_{\epsilon\gamma}^b p_\beta^{a-b} p_\gamma^b} \rightarrow \mathcal{B}_4 \\ \Psi_{\alpha\beta\gamma}[r(1-\kappa r\cos(\phi))] \sum_{b=-\infty}^{+\infty} (-ia\kappa \frac{B}{2A} P_\beta^{a-b} P_\gamma^b) := \sum_{b=-\infty}^{+\infty} -ia\kappa \frac{B}{2A} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \underline{\Psi_{\alpha\beta\gamma}[(1-\kappa r\cos(\phi))]p_\beta^{a-b} p_\gamma^b} \rightarrow \mathcal{B}_1 \end{cases} \quad (87)$$

Here, may be a little question of transform with P, think about  $N^{-1}$ , transform it as matrix we

could solve it:

$$\begin{aligned} & \because ia\kappa V^a = \frac{\partial P^a}{\partial r} + o(M^2) \\ & \int \int ia\kappa V^a \phi_\alpha dr d\theta = \int \int ia\kappa \phi_\alpha \phi_\beta dr d\theta V_\beta^a = \mathbf{I}_{\alpha\beta}^a V_\beta^a = \mathbf{V}_{\alpha\beta}^a p_\beta^a + \int \int o(M^2) r \phi_\alpha dr d\theta \\ & \because ia\kappa(1-\kappa r\cos\phi)V^a = (1-\kappa r\cos\phi) \frac{\partial P^a}{\partial r} + (1-\kappa r\cos\phi)o(M^2) \\ & \int \int ia\kappa(1-\kappa r\cos\phi)V^a \phi_\alpha dr d\theta = \mathbf{N}_{\alpha\beta}^a V_\beta^a = \int \int (1-\kappa r\cos\phi) \frac{\partial P^a}{\partial r} r \phi_\alpha dr d\theta + \int \int (1-\kappa r\cos\phi)o(M^2) dr d\theta \\ & \therefore (\mathbf{N}^{-1})_{\beta\alpha}^a \int \int (1-\kappa r\cos\phi) \frac{\partial P^a}{\partial r} r \phi_\alpha dr d\theta = (\mathbf{I}^{-1})_{\beta\alpha}^a \mathbf{I}_{\alpha\beta}^a V_\beta^a \\ & (\mathbf{N}^{-1})_{\beta\alpha}^a \int \int (1-\kappa r\cos\phi)o(M^2) dr d\theta = (\mathbf{I}^{-1})_{\beta\alpha}^a \int \int o(M^2) r \phi_\alpha dr d\theta \end{aligned} \quad (88)$$

## B. momentum equation I

$$\begin{aligned}
& \frac{d}{ds} P_\beta^a \widehat{\delta}_{\alpha\beta} - i a \kappa \Psi_{\alpha\beta} [r(1 - \kappa r \cos\phi)] U_\beta^a - \int_0^{2\pi} h h' [\psi_\beta \psi_\alpha]_{r=h} d\theta P_\beta^a - \Psi_{\{\alpha\}\beta} [r] P_\beta^a \\
&= \sum_{b=-\infty}^{+\infty} \underline{\text{term D1}} : \Psi_{\{\alpha\}\beta\gamma} [r] U_\beta^{a-b} U_\gamma^a + \Psi_{[\alpha]\beta\gamma} [r(1 - \kappa r \cos\phi)] \underline{\underline{V_\beta^{a-b} U_\gamma^a}} + \Psi_{(\alpha)\beta\gamma} [(1 - \kappa r \cos\phi)] \underline{\underline{W_\beta^{a-b} U_\gamma^a}} \\
&\quad + \underline{\text{term}(D1 + \mathcal{P}1)} : i(a) \kappa \Psi_{\alpha\beta\gamma} [r(1 - \kappa r \cos\phi)] P_\beta^{a-b} U_\gamma^b \\
&\quad + \underline{\text{term D1}} : -\frac{\partial}{\partial s} (\Psi_{\alpha\beta\gamma} [r]) U_\beta^{a-b} U_\gamma^b - \Psi_{\alpha\beta\gamma} [r] \left( \frac{dU_\beta^{a-b}}{ds} U_\gamma^b + \frac{dU_\gamma^b}{ds} U_\beta^{a-b} \right) \\
&\quad + \underline{\text{term X1}} : \Psi_{\alpha\beta\gamma} [r \cos\phi] U_\beta^{a-b} \underline{\underline{V_\gamma^b}} - \kappa \Psi_{\alpha\beta\gamma} [r \sin\phi] U_\beta^{a-b} \underline{\underline{W_\gamma^b}}
\end{aligned} \tag{89}$$

Transform:

$$\begin{aligned}
& \frac{d}{ds} P_\beta^a \widehat{\delta}_{\alpha\beta} := \widehat{\mathbf{I}_{\alpha\beta}^a} p_\beta'^a \\
& -ia \kappa \Psi_{\alpha\beta} [r(1 - \kappa r \cos\phi)] U_\beta^a := -\underline{\underline{ia \kappa \Psi_{\alpha\beta} [r(1 - \kappa \cos(\phi))]} u_\beta^a} \rightarrow -\mathcal{N} \\
& - \int_0^{2\pi} h h' [\psi_\beta \psi_\alpha]_{r=h} d\theta P_\beta^a - \Psi_{\{\alpha\}\beta} [r] P_\beta^a := -\underline{\underline{\int_0^{2\pi} h h' [\psi_\beta \psi_\alpha]_{r=h} d\theta p_\beta^a}} - \underline{\underline{\Psi_{\{\alpha\}\beta} [r] p_\beta^a}} = \underline{\underline{\Psi_{\{\alpha\}\beta} [r] p_\beta^a}} \rightarrow -\mathcal{H} \\
& \Psi_{\{\alpha\}\beta\gamma} [r] \sum_{b=-\infty}^{+\infty} U_\beta^{a-b} U_\gamma^a := \sum_{b=-\infty}^{+\infty} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \underline{\underline{\Psi_{\{\alpha\}\beta\gamma} [r] u_\beta^{a-b} u_\gamma^b}} \rightarrow \mathcal{D}_4 \\
& \Psi_{[\alpha]\beta\gamma} [r(1 - \kappa r \cos\phi)] \sum_{b=-\infty}^{+\infty} \underline{\underline{V_\beta^{a-b} U_\gamma^a}} := \sum_{b=-\infty}^{+\infty} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \sum_{\delta,\epsilon=0}^{\infty} \underline{\underline{\Psi_{[\alpha]\delta\epsilon} [r(1 - \kappa r \cos\phi)] \mathbf{V}_{\delta\beta}^{a-b} \mathbf{I}_{\epsilon\gamma}^b p_\beta^{a-b} u_\gamma^b}} \rightarrow \mathcal{C}_4 \\
& \Psi_{(\alpha)\beta\gamma} [(1 - \kappa r \cos\phi)] \sum_{b=-\infty}^{+\infty} \underline{\underline{W_\beta^{a-b} U_\gamma^a}} := \sum_{b=-\infty}^{+\infty} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \sum_{\delta,\epsilon=0}^{\infty} \underline{\underline{\Psi_{(\alpha)\delta\epsilon} [(1 - \kappa r \cos\phi)] \mathbf{W}_{\delta\beta}^{a-b} \mathbf{I}_{\epsilon\gamma}^b p_\beta^{a-b} u_\gamma^b}} \rightarrow \mathcal{C}_5 \\
& ia \kappa \Psi_{\alpha\beta\gamma} [r(1 - \kappa r \cos\phi)] \sum_{b=-\infty}^{+\infty} P_\beta^{a-b} U_\gamma^b := ia \kappa \sum_{b=-\infty}^{+\infty} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \underline{\underline{\Psi_{\alpha\beta\gamma} [r(1 - \kappa r \cos\phi)] p_\beta^{a-b} u_\gamma^b}} \rightarrow \mathcal{C}_3 \\
& -\frac{\partial}{\partial s} (\Psi_{\alpha\beta\gamma} [r]) \sum_{b=-\infty}^{+\infty} U_\beta^{a-b} U_\gamma^b := \sum_{b=-\infty}^{+\infty} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} -\frac{\partial}{\partial s} (\Psi_{\alpha\beta\gamma} [r]) u_\beta^{a-b} u_\gamma^b \rightarrow \mathcal{D}_1 \\
& -\Psi_{\alpha\beta\gamma} [r] \sum_{b=-\infty}^{+\infty} \left( \frac{dU_\beta^{a-b}}{ds} U_\gamma^b + \frac{dU_\gamma^b}{ds} U_\beta^{a-b} \right) := \Psi_{\alpha\beta\gamma} [r] \sum_{b=-\infty}^{+\infty} \left( [(Mp)_\beta^{a-b} + (Gu)_\beta^{a-b}] U_\gamma^b + [(Mp)_\gamma^b + (Gu)_\gamma^b] U_\beta^{a-b} \right) \\
& = \sum_{b=-\infty}^{+\infty} (\Psi_{\alpha\beta\gamma} [r] [\mathbf{G}, \mathbf{I}] + \Psi_{\alpha\beta\gamma} [r] [\mathbf{I}, \mathbf{G}]) u_\beta^{a-b} u_\gamma^b \rightarrow \mathcal{D}_{2,3} \\
& + \sum_{b=-\infty}^{+\infty} (\Psi_{\alpha\beta\gamma} [r] [\mathbf{M}, \mathbf{I}] + \Psi_{\alpha\beta\gamma} [r] [\mathbf{I}, \mathbf{M}]) u_\beta^{a-b} p_\gamma^b \rightarrow \mathcal{C}_{1,2} \\
& \Psi_{\alpha\beta\gamma} [r \cos\phi] U_\beta^{a-b} \underline{\underline{V_\gamma^b}} - \kappa \Psi_{\alpha\beta\gamma} [r \sin\phi] U_\beta^{a-b} \underline{\underline{W_\gamma^b}} := \{ \sum_{b=-\infty}^{+\infty} \Psi_{\alpha\beta\gamma} [r \cos\phi] [\mathbf{I}, \mathbf{V}] - \kappa \Psi_{\alpha\beta\gamma} [r \sin\phi] [\mathbf{I}, \mathbf{W}] \} u_\gamma^{a-b} p_\beta^b \rightarrow \mathcal{C}_{6,7}
\end{aligned} \tag{90}$$

Here, little transform easy to be proved:

$$\begin{aligned}
\Psi_{[\alpha]\beta\gamma}[r(1 - \kappa r \cos\phi)] \sum_{b=-\infty}^{+\infty} \underline{\underline{V_\beta^{a-b}}} U_\gamma^a &:= \sum_{b=-\infty}^{+\infty} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \sum_{\delta,\epsilon=0}^{\infty} \Psi_{[\alpha]\delta\epsilon}[r(1 - \kappa r \cos\phi)] \underline{\underline{\mathbf{V}_{\delta\beta}^{a-b} \mathbf{I}_{\epsilon\gamma}^b p_\beta^{a-b} u_\gamma^b}} \rightarrow \mathcal{C}_4 \\
\therefore \Psi_{[\alpha]\beta\gamma}[r(1 - \kappa r \cos\phi)][I, V][p_\beta^{a-b}, u_\gamma^b] &= \Psi_{[\alpha]\beta\gamma}[r(1 - \kappa r \cos\phi)][V, I][u_\beta^{a-b}, p_\gamma^b] \\
\Psi_{(\alpha)\beta\gamma}[(1 - \kappa r \cos\phi)] \sum_{b=-\infty}^{+\infty} \underline{\underline{W_\beta^{a-b}}} U_\gamma^a &:= \sum_{b=-\infty}^{+\infty} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \sum_{\delta,\epsilon=0}^{\infty} \Psi_{(\alpha)\delta\epsilon}[(1 - \kappa r \cos\phi)] \underline{\underline{\mathbf{W}_{\delta\beta}^{a-b} \mathbf{I}_{\epsilon\gamma}^b p_\beta^{a-b} u_\gamma^b}} \rightarrow \mathcal{C}_5 \\
\therefore \Psi_{(\alpha)\beta\gamma}[1 - \kappa r \cos\phi][I, W][p_\beta^{a-b}, u_\gamma^b] &= \Psi_{[\alpha]\beta\gamma}[1 - \kappa r \cos\phi][W, I][u_\beta^{a-b}, p_\gamma^b] \\
ia\kappa \Psi_{\alpha\beta\gamma}[r(1 - \kappa r \cos\phi)] \sum_{b=-\infty}^{+\infty} P_\beta^{a-b} U_\gamma^b &:= ia\kappa \sum_{b=-\infty}^{+\infty} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \underline{\underline{\Psi_{\alpha\beta\gamma}[r(1 - \kappa r \cos\phi)] p_\beta^{a-b} u_\gamma^b}} \rightarrow \mathcal{C}_3 \\
\therefore ia\kappa \Psi_{\alpha\beta\gamma}[r(1 - \kappa r \cos\phi)][I, I][p_\beta^{a-b}, u_\gamma^b] &= ia\kappa \Psi_{\alpha\beta\gamma}[r(1 - \kappa r \cos\phi)][I, I][u_\beta^{a-b}, p_\gamma^b]
\end{aligned} \tag{91}$$

### C. momentum equation II

$$\begin{aligned}
&-ia\kappa \Psi_{\alpha\beta}[r(1 - \kappa r \cos\phi)] V_\beta^a \\
&+ \int_0^{2\pi} [\psi_\alpha \psi_\beta r(1 - \kappa r \cos\phi)]_0^h d\theta P_\beta^a - \Psi_{[\alpha]\beta}[r(1 - \kappa r \cos\phi)] P_\beta^a - \Psi_{\alpha\beta}[1 - 2\kappa r \cos\phi] P_\beta^a \\
&= \underline{\text{term D2}} : \Psi_{\{\alpha\}\beta\gamma}[r] U_\beta^{a-b} V_\gamma^a + \Psi_{[\alpha]\beta\gamma}[r(1 - \kappa r \cos\phi)] V_\beta^{a-b} V_\gamma^a + \Psi_{(\alpha)\beta\gamma}[(1 - \kappa r \cos\phi)] W_\beta^{a-b} V_\gamma^a \\
&\quad + \underline{\text{term}(D2 + P2)} : i(a) \kappa \Psi_{\alpha\beta\gamma}[r(1 - \kappa r \cos\phi)] P_\beta^{a-b} V_\gamma^b \\
&\quad + \underline{\text{term D2}} : -\frac{\partial}{\partial s} (\Psi_{\alpha\beta\gamma}[r]) U_\beta^{a-b} V_\gamma^b - \Psi_{\alpha\beta\gamma}[r] \left( \frac{dU_\beta^{a-b}}{ds} V_\gamma^b + \frac{dV_\gamma^b}{ds} U_\beta^{a-b} \right) \\
&\quad + \underline{\text{term X2}} : \Psi_{\alpha\beta\gamma}[1 - \kappa r \cos\phi] W_\beta^{a-b} W_\gamma^b - \kappa \Psi_{\alpha\beta\gamma}[r \cos\phi] U_\beta^{a-b} U_\gamma^b
\end{aligned} \tag{92}$$

With:

$$\begin{aligned}
\frac{d}{ds} V_\beta^a \widehat{\delta}_{\alpha\beta} &= \left\{ \int_0^{2\pi} \frac{h'^2}{1 - \kappa h \cos\phi} [r \psi_\beta \psi_\alpha]_0^h d\theta \right. \\
&\quad \left. + \int_0^{2\pi} [r(1 - \kappa r \cos\phi) \psi_\beta \psi_\alpha]_0^h d\theta - \Psi_{[\alpha]\beta}[r(1 - \kappa r \cos\phi)] - \Psi_{\alpha\beta}[(1 - \kappa r \cos\phi)] \right\} U_\beta^a \\
&\quad - G_{\alpha\beta}^a V_\beta^a
\end{aligned} \tag{93}$$

Transform:

$$\begin{aligned}
& -ia\kappa\Psi_{\alpha\beta}[r(1-\kappa r\cos\phi)]V_\beta^a := -N_{\alpha\beta}^a V_\beta^a \\
& \left\{ \int_0^{2\pi} [\psi_\alpha\psi_\beta r(1-\kappa r\cos\phi)]_0^h d\theta - \Psi_{[\alpha]\beta}[r(1-\kappa r\cos\phi)] - \Psi_{\alpha\beta}[1-2\kappa r\cos\phi] \right\} P_\beta^a \\
& \Psi_{\{\alpha\}\beta\gamma}[r]U_\beta^{a-b}V_\gamma^a := \sum_{b=-\infty}^{+\infty} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \underline{\Psi_{\{\alpha\}\delta\epsilon}[r]\mathbf{I}_{\delta\beta}^{a-b}\mathbf{V}_{\epsilon\gamma}^b u_\beta^{a-b} p_\gamma^b} = \sum_{b=-\infty}^{+\infty} \Psi_{\{\alpha\}\beta\gamma}[r][\mathbf{I}, \mathbf{V}]u_\beta^{a-b} p_\gamma^b \rightarrow \varepsilon_{1.4} \\
& \Psi_{[\alpha]\beta\gamma}[r(1-\kappa r\cos\phi)]V_\beta^{a-b}V_\gamma^a := \sum_{b=-\infty}^{+\infty} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \underline{\Psi_{[\alpha]\delta\epsilon}[r(1-\kappa r\cos\phi)]\mathbf{V}_{\delta\beta}^{a-b}\mathbf{V}_{\epsilon\gamma}^b p_\beta^{a-b} p_\gamma^b} \rightarrow \mathcal{B}_{5.3} \\
& \Psi_{(\alpha)\beta\gamma}[(1-\kappa r\cos\phi)]W_\beta^{a-b}V_\gamma^a := \sum_{b=-\infty}^{+\infty} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \underline{\Psi_{(\alpha)\delta\epsilon}[1-\kappa r\cos\phi]\mathbf{W}_{\delta\beta}^{a-b}\mathbf{V}_{\epsilon\gamma}^b p_\beta^{a-b} p_\gamma^b} \rightarrow \mathcal{B}_{5.4} \\
& i(a)\kappa\Psi_{\alpha\beta\gamma}[r(1-\kappa r\cos\phi)]P_\beta^{a-b}V_\gamma^b := \sum_{b=-\infty}^{+\infty} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \underline{i\kappa\Psi_{\alpha\delta\epsilon}[r(1-\kappa r\cos\phi)]\mathbf{I}_{\delta\beta}^{a-b}\mathbf{V}_{\epsilon\gamma}^b p_\beta^{a-b} p_\gamma^b} \rightarrow \mathcal{B}_{5.2} \\
& -\frac{\partial}{\partial s}(\Psi_{\alpha\beta\gamma}[r])U_\beta^{a-b}V_\gamma^b := \sum_{b=-\infty}^{+\infty} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} -\frac{\partial}{\partial s}(\Psi_{\alpha\delta\epsilon}[r])\mathbf{I}_{\delta\beta}^{a-b}\mathbf{V}_{\epsilon\gamma}^b u_\beta^{a-b} p_\gamma^b \rightarrow \varepsilon_{1.1} \\
& -\Psi_{\alpha\beta\gamma}[r] \sum_{b=-\infty}^{+\infty} (\widehat{\mathbf{I}^{a-b}}^{-1}[-(\mathbf{M}p)_\beta^{a-b} - (\mathbf{G}u)_\beta^{a-b}]V_\gamma^b \\
& := -\Psi_{\alpha\beta\gamma}[r] \sum_{b=-\infty}^{+\infty} (\widehat{\mathbf{I}^{a-b}}^{-1}[-(\mathbf{M}p)_\beta^{a-b} - (\mathbf{G}u)_\beta^{a-b}]V_\gamma^b \\
& \quad + \widehat{\mathbf{I}_{\gamma\alpha}^b}^{-1} \left\{ \int_0^{2\pi} \frac{h'^2}{1-\kappa h\cos\phi} [r\psi_\beta\psi_\alpha]_0^h d\theta \right. \\
& \quad \left. + \int_0^{2\pi} [r(1-\kappa r\cos\phi)\psi_\beta\psi_\alpha]_0^h d\theta - \Psi_{[\alpha]\beta}[r(1-\kappa r\cos\phi)] - \Psi_{\alpha\beta}[(1-\kappa r\cos\phi)] \right\} U_\beta^b U_\beta^{a-b} \\
& \quad - \sum_{b=-\infty}^{+\infty} \Psi_{\alpha\beta\gamma}[r] \left\{ \widehat{\mathbf{I}}^{-1} \left( \int_0^{2\pi} \frac{h'^2}{1-\kappa h\cos\phi} [r\psi_\beta\psi_\alpha]_0^h d\theta \right) \right. \\
& \quad \left. + \sum_{b=-\infty}^{+\infty} \Psi_{\alpha\beta\gamma}[r][\widehat{\mathbf{I}}, \widehat{\mathbf{I}}^{-1}\mathbf{GV}]u_\beta^{a-b} p_\gamma^b \right\} \rightarrow \varepsilon_{1.2} \\
& \quad + \sum_{b=-\infty}^{+\infty} (\Psi_{\alpha\beta\gamma}[r]\widehat{\mathbf{I}}^{-1}\mathbf{M}, \mathbf{V}))p_\beta^{a-b} p_\gamma^b \rightarrow \varepsilon_{1.2} \\
& \quad + \sum_{b=-\infty}^{+\infty} (\Psi_{\alpha\beta\gamma}[r]\widehat{\mathbf{I}}^{-1}\mathbf{M}, \mathbf{V}))p_\beta^{a-b} p_\gamma^b \rightarrow \mathcal{B}_{5.1} \\
& \quad - \sum_{b=-\infty}^{+\infty} \Psi_{\alpha\beta\gamma}[r] \left\{ \widehat{\mathbf{I}}^{-1} \left( \int_0^{2\pi} \frac{h'^2}{1-\kappa h\cos\phi} [r\psi_\beta\psi_\alpha]_0^h d\theta \right) \right. \\
& \quad \left. + \sum_{b=-\infty}^{+\infty} \Psi_{\alpha\beta\gamma}[r][\mathbf{I}, \widehat{\mathbf{I}}^{-1}\mathbf{GV}]u_\beta^{a-b} p_\gamma^b \right\} \rightarrow \varepsilon_{1.3} \\
& \quad + \sum_{b=-\infty}^{+\infty} \Psi_{\alpha\beta\gamma}[r][\mathbf{I}, \widehat{\mathbf{I}}^{-1}\mathbf{GV}]u_\beta^{a-b} p_\gamma^b \rightarrow \varepsilon_{1.3} \\
& \Psi_{\alpha\beta\gamma}[1-\kappa r\cos\phi]W_\beta^{a-b}W_\gamma^b := \sum_{b=-\infty}^{+\infty} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \underline{\Psi_{\alpha\delta\epsilon}[1-\kappa r\cos\phi]\mathbf{W}_{\delta\beta}^{a-b}\mathbf{W}_{\epsilon\gamma}^b p_\beta^{a-b} p_\gamma^b} \rightarrow \mathcal{B}_{5.5} \\
& -\kappa\Psi_{\alpha\beta\gamma}[r\cos\phi]U_\beta^{a-b}U_\gamma^b := \sum_{b=-\infty}^{+\infty} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} -\kappa\Psi_{\alpha\beta\gamma}[r\cos\phi]u_\beta^{a-b} u_\gamma^b \rightarrow \mathcal{A}_{2.5}
\end{aligned} \tag{94}$$

#### D. momentum equation III

$$\begin{aligned}
& -ia\kappa\Psi_{\alpha\beta}[r(1-\kappa r\cos\phi)]W_\beta^a \\
& -\Psi_{(\alpha)\beta}[(1-\kappa r\cos\phi)]P_\beta^a - \kappa\Psi_{\alpha\beta}[r\sin\phi]P_\beta^a \\
= & \underline{\text{termD2}} : \Psi_{\{\alpha\}\beta\gamma}[r]U_\beta^{a-b}W_\gamma^a + \Psi_{[\alpha]\beta\gamma}[r(1-\kappa r\cos\phi)]V_\beta^{a-b}W_\gamma^a + \Psi_{(\alpha)\beta\gamma}[(1-\kappa r\cos\phi)]W_\beta^{a-b}W_\gamma^a \\
& + \underline{\text{term}(D2 + \mathcal{P}2)} : i(a)\kappa\Psi_{\alpha\beta\gamma}[r(1-\kappa r\cos\phi)]P_\beta^{a-b}W_\gamma^a \\
& + \underline{\text{termD2}} : -\frac{\partial}{\partial s}(\Psi_{\alpha\beta\gamma}[r])U_\beta^{a-b}V_\gamma^b - \Psi_{\alpha\beta\gamma}[r](\frac{dU_\beta^{a-b}}{ds}W_\gamma^b + \frac{dW_\gamma^b}{ds}U_\beta^{a-b}) \\
& + \underline{\text{termX2}} : \kappa\Psi_{\alpha\beta\gamma}[r\sin\phi]U_\beta^{a-b}U_\gamma^b - \Psi_{\alpha\beta\gamma}[1-\kappa r\cos\phi]W_\beta^{a-b}V_\gamma^b
\end{aligned} \tag{95}$$

With:

---


$$\frac{d}{ds}W_\beta^a\hat{\delta}_{\alpha\beta} = \int_0^{2\pi} \frac{dh(s)}{ds} [\psi_\beta r\psi_\alpha]_0^h d\theta W_\beta^a + \Psi_{\{\alpha\}\beta}[r]W_\beta^a - \Psi_{(\alpha)\beta}[1-\kappa r\cos\phi]U_\beta^a = -\Psi_{\alpha\{\beta\}}[r]W_\beta^a - \Psi_{(\alpha)\beta}[1-\kappa r\cos\phi]U_\beta^a$$


---

Transform:

$$\begin{aligned}
& -ia\kappa\Psi_{\alpha\beta}[r(1-\kappa r\cos\phi)]W_\beta^a := -N_{\alpha\beta}^a W_\beta^a \\
& -\Psi_{(\alpha)\beta}[(1-\kappa r\cos\phi)]p_\beta^a - \kappa\Psi_{\alpha\beta}[r\sin\phi]p_\beta^a \\
\Psi_{\{\alpha\}\beta\gamma}[r]U_\beta^{a-b}W_\gamma^a &:= \sum_{b=-\infty}^{+\infty} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \underline{\Psi_{\{\alpha\}\delta\epsilon}[r]\mathbf{I}_{\delta\beta}^{a-b}\mathbf{W}_{\epsilon\gamma}^b} u_\beta^{a-b} p_\gamma^b = \sum_{b=-\infty}^{+\infty} \Psi_{\{\alpha\}\beta\gamma}[r][\mathbf{I}, \mathbf{W}]u_\beta^{a-b} p_\gamma^b \rightarrow \varepsilon_{2.4} \\
\Psi_{[\alpha]\beta\gamma}[r(1-\kappa r\cos\phi)]V_\beta^{a-b}W_\gamma^a &:= \sum_{b=-\infty}^{+\infty} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \underline{\Psi_{[\alpha]\delta\epsilon}[r(1-\kappa r\cos\phi)]\mathbf{V}_{\delta\beta}^{a-b}\mathbf{W}_{\epsilon\gamma}^b} p_\beta^{a-b} p_\gamma^b \rightarrow \mathcal{B}_{6.3} \\
\Psi_{(\alpha)\beta\gamma}[(1-\kappa r\cos\phi)]W_\beta^{a-b}W_\gamma^a &:= \sum_{b=-\infty}^{+\infty} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \underline{\Psi_{(\alpha)\delta\epsilon}[1-\kappa r\cos\phi]\mathbf{W}_{\delta\beta}^{a-b}\mathbf{W}_{\epsilon\gamma}^b} p_\beta^{a-b} p_\gamma^b \rightarrow \mathcal{B}_{6.4} \\
i(a)\kappa\Psi_{\alpha\beta\gamma}[r(1-\kappa r\cos\phi)]P_\beta^{a-b}W_\gamma^b &:= \sum_{b=-\infty}^{+\infty} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \underline{i\kappa\Psi_{\alpha\delta\epsilon}[r(1-\kappa r\cos\phi)]\mathbf{I}_{\delta\beta}^{a-b}\mathbf{W}_{\epsilon\gamma}^b} p_\beta^{a-b} p_\gamma^b \rightarrow \mathcal{B}_{6.2} \\
-\frac{\partial}{\partial s}(\Psi_{\alpha\beta\gamma}[r])U_\beta^{a-b}W_\gamma^b &:= \sum_{b=-\infty}^{+\infty} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} -\frac{\partial}{\partial s}(\Psi_{\alpha\delta\epsilon}[r])\mathbf{I}_{\delta\beta}^{a-b}\mathbf{W}_{\epsilon\gamma}^b u_\beta^{a-b} p_\gamma^b \rightarrow \varepsilon_{2.1} \\
& -\Psi_{\alpha\beta\gamma}[r] \sum_{b=-\infty}^{+\infty} \left( \frac{dU_\beta^{a-b}}{ds} W_\gamma^b + \frac{dW_\gamma^b}{ds} U_\beta^{a-b} \right) \\
& := -\Psi_{\alpha\beta\gamma}[r] \sum_{b=-\infty}^{+\infty} (\widehat{\mathbf{I}}^{-1}[-(\mathbf{M}p)_\beta^{a-b} - (\mathbf{G}u)_\beta^{a-b}]W_\gamma^b - \{\mathbf{H}W_\beta^a - \Psi_{(\alpha)\beta}[1-\kappa r\cos\phi]U_\beta^a\}U_\beta^{a-b} \\
& = \sum_{b=-\infty}^{+\infty} (\Psi_{\alpha\beta\gamma}[r][\widehat{\mathbf{I}}^{-1}\mathbf{G}, \mathbf{W}])u_\beta^{a-b} p_\gamma^b \rightarrow \varepsilon_{2.2} \\
& + \sum_{b=-\infty}^{+\infty} (\Psi_{\alpha\beta\gamma}[r][\widehat{\mathbf{I}}^{-1}\mathbf{M}, \mathbf{W}])p_\beta^{a-b} p_\gamma^b \rightarrow \mathcal{B}_{6.1} \\
& - \sum_{b=-\infty}^{+\infty} \Psi_{\alpha\beta\gamma}[r][\mathbf{I}, \widehat{\mathbf{I}}^{-1}\mathbf{H}\mathbf{W}]u_\beta^{a-b} p_\gamma^b \rightarrow \varepsilon_{2.3} \\
& + \sum_{b=-\infty}^{+\infty} \Psi_{\alpha\beta\gamma}[r](\widehat{\mathbf{I}}^{-1}\Psi_{(\alpha)\beta}[1-\kappa r\cos\phi])u_\beta^a u_\beta^{a-b} \rightarrow \mathcal{A}_{3.1} \\
& \kappa\Psi_{\alpha\beta\gamma}[r\sin\phi]U_\beta^{a-b}U_\gamma^b := \sum_{b=-\infty}^{+\infty} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \underline{\kappa\Psi_{\alpha\beta\gamma}[r\sin\phi]} u_\beta^{a-b} u_\gamma^b \rightarrow \mathcal{A}_{3.2} \\
& -\Psi_{\alpha\beta\gamma}[1-\kappa r\cos\phi]W_\beta^{a-b}V_\gamma^b := \sum_{b=-\infty}^{+\infty} \sum_{\beta=0}^{+\infty} \sum_{\gamma=0}^{+\infty} \underline{-\Psi_{\alpha\delta\epsilon}[1-\kappa r\cos\phi]\mathbf{W}_{\delta\beta}^{a-b}\mathbf{V}_{\epsilon\gamma}^b} p_\beta^{a-b} p_\gamma^b \rightarrow \mathcal{B}_{6.5}
\end{aligned} \tag{97}$$

Finally, we obtain two equations involving just pressure and longitudinal velocity modes. Here,  $\mathbf{M}, \mathbf{N}, \mathcal{A}, \mathcal{B}$ , and  $\mathcal{C}$  encoding the curvature of the duct. The terms  $\mathbf{G}, \mathbf{H}, \mathbf{D}$  encode the variation in duct diameter as well as the torsion. The term  $\varepsilon$  encodes variation of diameter and the torsion together with curvature if either of the first two are present.

$$\begin{aligned}
\widehat{\mathbf{I}_{\alpha\beta}^a} u'_\beta + \mathbf{M}_{\alpha\beta}^a p_\beta^a + \mathbf{G}_{\alpha\beta}^a u_\beta^a &= \mathcal{A}_{\alpha\beta\gamma}^{ab}[u_\beta^{a-b}, u_\gamma^b] + \mathcal{B}_{\alpha\beta\gamma}^{ab}[p_\beta^{a-b}, p_\gamma^b] + \varepsilon_{\alpha\beta\gamma}^{ab}[u_\beta^{a-b}, p_\gamma^b] \\
\widehat{\mathbf{I}_{\alpha\beta}^a} p'_\beta - \mathbf{N}_{\alpha\beta}^a u_\beta^a - \mathbf{H}_{\alpha\beta}^a p_\beta^a &= \mathcal{C}_{\alpha\beta\gamma}^{ab}[u_\beta^{a-b}, p_\gamma^b] + \mathcal{D}_{\alpha\beta\gamma}^{ab}[u_\beta^{a-b}, u_\gamma^b]
\end{aligned} \tag{98}$$

## VI. Introduce the admittance matrix

Due to the present of evanescent modes these equations are numerically unstable and cannot be integrated directly. Define a relation between the pressure and velocity in terms of the admittance. When solving for pressure, it is easier to work with the admittance rather than the impedance Z ( $Y = Z^{-1}$ ), to avoid inverting large matrices in the work that will follow.

The following relationship is defined:

$$\widehat{\mathbf{I}_{\alpha\beta}^a} u_\beta^a = Y_{\alpha\beta}^a p_\beta^a + \mathcal{Y}_{\alpha\beta\gamma}^{ab}[p_\beta^{a-b}, p_\gamma^b] \tag{99}$$

where  $Y = Y(s)$  is the linear part of the admittance and  $\mathcal{Y} = \mathcal{Y}(s)$  is the second order nonlinear correction to the admittance, henceforth referred to as the nonlinear admittance term. We differentiate it:

$$\widehat{\mathbf{I}_{\alpha\beta}^a} u' = Y'_{\alpha\beta}^a p_\beta^a + Y_{\alpha\beta}^a p'_\beta + \mathcal{Y}'_{\alpha\beta\gamma}^{ab}[p_\beta^{a-b}, p_\gamma^b] + \mathcal{Y}_{\alpha\beta\gamma}^{ab}[p_\beta'^{a-b}, p_\gamma^b] + \mathcal{Y}_{\alpha\beta\gamma}^{ab}[p_\beta^{a-b}, p_\gamma'^b] \tag{100}$$

Substitute in eq98 of  $u'$ ,

$$\begin{aligned}
&- \mathbf{M}_{\alpha\beta}^a p_\beta^a - \mathbf{G}_{\alpha\beta}^a u_\beta^a + \mathcal{A}_{\alpha\beta\gamma}^{ab}[u_\beta^{a-b}, u_\gamma^b] + \mathcal{B}_{\alpha\beta\gamma}^{ab}[p_\beta^{a-b}, p_\gamma^b] + \varepsilon_{\alpha\beta\gamma}^{ab}[u_\beta^{a-b}, p_\gamma^b] \\
&= Y'_{\alpha\beta}^a p_\beta^a + Y_{\alpha\beta}^a p'_\beta + \mathcal{Y}'_{\alpha\beta\gamma}^{ab}[p_\beta^{a-b}, p_\gamma^b] + \mathcal{Y}_{\alpha\beta\gamma}^{ab}[p_\beta'^{a-b}, p_\gamma^b] + \mathcal{Y}_{\alpha\beta\gamma}^{ab}[p_\beta^{a-b}, p_\gamma'^b]
\end{aligned} \tag{101}$$

Then,  $p'$

$$\begin{aligned}
&- \mathbf{M}_{\alpha\beta}^a p_\beta^a - \mathbf{G}_{\alpha\beta}^a u_\beta^a + \mathcal{A}_{\alpha\beta\gamma}^{ab}[u_\beta^{a-b}, u_\gamma^b] + \mathcal{B}_{\alpha\beta\gamma}^{ab}[p_\beta^{a-b}, p_\gamma^b] + \varepsilon_{\alpha\beta\gamma}^{ab}[u_\beta^{a-b}, p_\gamma^b] \\
&= Y'_{\alpha\beta}^a p_\beta^a \\
&+ Y_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} (\mathbf{N}_{\alpha\beta}^a u_\beta^a + \mathbf{H}_{\alpha\beta}^a p_\beta^a + \mathcal{C}_{\alpha\beta\gamma}^{ab}[u_\beta^{a-b}, p_\gamma^b] + \mathcal{D}_{\alpha\beta\gamma}^{ab}[u_\beta^{a-b}, u_\gamma^b]) \\
&+ \mathcal{Y}'_{\alpha\beta\gamma}^{ab}[p_\beta^{a-b}, p_\gamma^b] \\
&+ \mathcal{Y}_{\alpha\beta\gamma}^{ab}[\mathbf{N}_{\beta\delta}^{a-b} u_\delta^{a-b} + \mathbf{H}_{\beta\delta}^{a-b} p_\delta^{a-b}, p_\gamma^b] \\
&+ \mathcal{Y}_{\alpha\beta\gamma}^{ab}[p_\beta^{a-b}, \mathbf{N}_{\gamma\delta}^b u_\delta^b + \mathbf{H}_{\gamma\delta}^b p_\delta^b]
\end{aligned} \tag{102}$$

Use eq99 to express  $u$  in terms of  $p$ ,  $\widehat{\mathbf{I}}_{\alpha\beta}^a u_\beta^a = Y_{\alpha\beta}^a p_\beta^a + \mathcal{Y}_{\alpha\beta\gamma}^{ab}[p_\beta^{a-b}, p_\gamma^b]$

$$\begin{aligned}
& -\mathbf{M}_{\alpha\beta}^a p_\beta^a - \mathbf{G}_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^a{}^{-1} Y_{\alpha\beta}^a p_\beta^a - \mathbf{G}_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^a{}^{-1} \mathcal{Y}_{\alpha\beta\gamma}^{ab}[p_\beta^{a-b}, p_\gamma^b] \\
& + \mathcal{A}_{\alpha\beta\gamma}^{ab} [\widehat{\mathbf{I}}_{\beta\alpha}^{a-b}{}^{-1} Y_{\alpha\beta}^{\pm a-b} p_\beta^{a-b}, \widehat{\mathbf{I}}_{\gamma\alpha}^{a-b}{}^{-1} Y_{\alpha\gamma}^{\pm b} p_\gamma^b] + \mathcal{B}_{\alpha\beta\gamma}^{ab}[p_\beta^{a-b}, p_\gamma^b] + \varepsilon_{\alpha\beta\gamma}^{ab} [\widehat{\mathbf{I}}_{\beta\alpha}^{a-b}{}^{-1} Y_{\alpha\beta}^{\pm a-b} p_\beta^{a-b}, p_\gamma^b] \\
& = Y'_{\alpha\beta}^a p_\beta^a + Y_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^a{}^{-1} (\mathbf{N}_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^a{}^{-1} Y_{\alpha\beta}^a p_\beta^a + \mathbf{N}_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^a{}^{-1} \mathcal{Y}_{\alpha\beta\gamma}^{ab}[p_\beta^{a-b}, p_\gamma^b] + \mathbf{H}_{\alpha\beta}^a p_\beta^a) \\
& + \mathcal{C}_{\alpha\beta\gamma}^{ab} [\widehat{\mathbf{I}}_{\beta\alpha}^{a-b}{}^{-1} Y_{\alpha\beta}^{\pm a-b} p_\beta^{a-b}, p_\gamma^b] + \mathcal{D}_{\alpha\beta\gamma}^{ab} [\widehat{\mathbf{I}}_{\beta\alpha}^{a-b}{}^{-1} Y_{\alpha\beta}^{\pm a-b} p_\beta^{a-b}, \widehat{\mathbf{I}}_{\gamma\alpha}^{a-b}{}^{-1} Y_{\alpha\gamma}^{\pm b} p_\gamma^b] \\
& + \mathcal{Y}_{\alpha\beta\gamma}^{ab}[p_\beta^{a-b}, p_\gamma^b] + \mathcal{Y}_{\alpha\beta\gamma}^{ab} [\mathbf{N}_{\beta\delta}^a \widehat{\mathbf{I}}_{\delta\alpha}^{a-b}{}^{-1} Y_{\alpha\delta}^{a-b} p_\delta^{a-b} + \mathbf{H}_{\beta\delta}^{a-b} p_\delta^{a-b}, p_\gamma^b] \\
& + \mathcal{Y}_{\alpha\beta\gamma}^{ab}[p_\beta^{a-b}, \mathbf{N}_{\gamma\delta}^b \widehat{\mathbf{I}}_{\delta\alpha}^{b-a}{}^{-1} Y_{\alpha\delta}^b p_\delta^b + \mathbf{H}_{\gamma\delta}^b p_\delta^b]
\end{aligned} \tag{103}$$

This equation has two distinct orders of magnitude: terms linear in  $p$ , and terms quadratic in

p. We can equate linear terms and the quadratic terms separately to get two distinct equations:

$$\text{linear : } Y'_{\alpha\beta}^a p_\beta^a + Y_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^a{}^{-1} \mathbf{N}_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^a{}^{-1} Y_{\alpha\beta}^a p_\beta^a + \mathbf{M}_{\alpha\beta}^a p_\beta^a + Y_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^a{}^{-1} \mathbf{H}_{\alpha\beta}^a p_\beta^a + \mathbf{G}_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^a{}^{-1} Y_{\alpha\beta}^a p_\beta^a = 0, \tag{104}$$

$$\begin{aligned}
\text{quadratic : } & \mathcal{Y}_{\alpha\beta\gamma}^{ab}[p_\beta^{a-b}, p_\gamma^b] + \mathcal{Y}_{\alpha\beta\gamma}^{ab} [\mathbf{N}_{\beta\delta}^a \widehat{\mathbf{I}}_{\delta\alpha}^{a-b}{}^{-1} Y_{\alpha\delta}^{a-b} p_\delta^{a-b}, p_\gamma^b] + \mathcal{Y}_{\alpha\beta\gamma}^{ab}[p_\beta^{a-b}, \mathbf{N}_{\gamma\delta}^b \widehat{\mathbf{I}}_{\delta\alpha}^{b-a}{}^{-1} Y_{\alpha\delta}^b p_\delta^b] \\
& + Y_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^a{}^{-1} \mathbf{N}_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^a{}^{-1} \mathcal{Y}_{\alpha\beta\gamma}^{ab}[p_\beta^{a-b}, p_\gamma^b] + Y_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^a{}^{-1} \mathcal{C}_{\alpha\beta\gamma}^{ab} [\widehat{\mathbf{I}}_{\beta\alpha}^{a-b}{}^{-1} Y_{\alpha\beta}^{\pm a-b} p_\beta^{a-b}, p_\gamma^b] \\
& - \mathcal{A}_{\alpha\beta\gamma}^{ab} [\widehat{\mathbf{I}}_{\beta\alpha}^{a-b}{}^{-1} Y_{\alpha\beta}^{\pm a-b} p_\beta^{a-b}, \widehat{\mathbf{I}}_{\gamma\alpha}^{a-b}{}^{-1} Y_{\alpha\gamma}^{\pm b} p_\gamma^b] - \mathcal{B}_{\alpha\beta\gamma}^{ab}[p_\beta^{a-b}, p_\gamma^b] - \varepsilon_{\alpha\beta\gamma}^{ab} [\widehat{\mathbf{I}}_{\beta\alpha}^{a-b}{}^{-1} Y_{\alpha\beta}^{\pm a-b} p_\beta^{a-b}, p_\gamma^b] \\
& + \mathcal{Y}_{\alpha\beta\gamma}^{ab}[p_\beta^{a-b}, \mathbf{H}_{\gamma\delta}^b p_\delta^b] + \mathcal{Y}_{\alpha\beta\gamma}^{ab} [\mathbf{H}_{\beta\delta}^{a-b} p_\delta^{a-b}, p_\gamma^b] + \mathbf{G}_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^a{}^{-1} \mathcal{Y}_{\alpha\beta\gamma}^{ab}[p_\beta^{a-b}, p_\gamma^b] \\
& + Y_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^a{}^{-1} \mathcal{D}_{\alpha\beta\gamma}^{ab} [\widehat{\mathbf{I}}_{\beta\alpha}^{a-b}{}^{-1} Y_{\alpha\beta}^{\pm a-b} p_\beta^{a-b}, \widehat{\mathbf{I}}_{\gamma\alpha}^{a-b}{}^{-1} Y_{\alpha\gamma}^{\pm b} p_\gamma^b] = 0
\end{aligned} \tag{105}$$

As both of these equations hold true for a general  $p$  we can eliminate it to obtain an equation

for the linear part of the admittance and an equation for the nonlinear part of the admittance:

$$\text{linear} - 2D : Y'_{\alpha\beta}^a + Y_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^a{}^{-1} \mathbf{N}_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^a{}^{-1} Y_{\alpha\beta}^a + \mathbf{M}_{\alpha\beta}^a + Y_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^a{}^{-1} \mathbf{H}_{\alpha\beta}^a + \mathbf{G}_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^a{}^{-1} Y_{\alpha\beta}^a = 0, \tag{106}$$

For quadratic, with:

$$\begin{aligned} (\mathcal{A}[x, y])_{\alpha}^a &= (\mathcal{A}_{\alpha\beta\gamma}^{ab}[x_{\beta}^{a-b}, y_{\gamma}^b])_{\alpha}^a = \sum_{b=-\infty}^{\infty} \sum_{\beta, \gamma=0}^{\infty} A_{\alpha\beta\gamma}^{ab} x_{\beta}^{a-b} y_{\gamma}^b \\ (\mathcal{A}[X, Y])_{\alpha\beta\gamma}^{ab} &= (\mathcal{A}_{\alpha\delta\epsilon}^{ab}[X_{\delta\beta}^{a-b}, Y_{\epsilon\gamma}^b])_{\alpha\beta\gamma}^{ab} = \sum_{\delta, \epsilon=0}^{\infty} A_{\alpha\delta\epsilon}^{ab} X_{\delta\beta}^{a-b} Y_{\epsilon\gamma}^b \end{aligned}$$

$$\begin{aligned} \text{Thus, } \{(\mathcal{A}[X, Y])_{\alpha\beta\gamma}^{ab}[x, y]\}_{\alpha}^a &= \{(\mathcal{A}_{\alpha\delta\epsilon}^{ab}[X_{\delta\beta}^{a-b}, Y_{\epsilon\gamma}^b])_{\alpha\beta\gamma}^{ab}[x_{\beta}^{a-b}, y_{\gamma}^b]\}_{\alpha}^a \\ &= \sum_{b=-\infty}^{\infty} \sum_{\beta, \gamma=0}^{\infty} \left( \sum_{\delta, \epsilon=0}^{\infty} A_{\alpha\delta\epsilon}^{ab} X_{\delta\beta}^{a-b} Y_{\epsilon\gamma}^b \right) x_{\beta}^{a-b} y_{\gamma}^b \\ &= \sum_{b=-\infty}^{\infty} \left\{ \sum_{\delta, \epsilon=0}^{\infty} A_{\alpha\delta\epsilon}^{ab} \left( \sum_{\beta} X_{\delta\beta}^{a-b} x_{\beta}^{a-b} \right)_{\delta}^{a-b} \left( \sum_{\gamma=0}^{\infty} Y_{\epsilon\gamma}^b \right)_{\epsilon}^b \right\} = \mathcal{A}[Xx, Yy] \end{aligned} \quad (107)$$

We now can eliminate p, remaining 3-D tensors:

$$\begin{aligned} \text{quadratic : } &\mathcal{Y}'_{\alpha\beta\gamma}^{ab}[I_{\beta\delta}p_{\delta}^{a-b}, I_{\gamma\epsilon}p_{\epsilon}^b] + \mathcal{Y}_{\alpha\beta\gamma}^{ab}[\mathbf{N}_{\beta\delta}^{a-b}\widehat{\mathbf{I}_{\delta\alpha}^{a-b}}^{-1} Y_{\alpha\delta}^{a-b} p_{\delta}^{a-b}, I_{\gamma\epsilon}p_{\epsilon}^b] + \mathcal{Y}_{\alpha\beta\gamma}^{ab}[I_{\beta\delta}p_{\delta}^{a-b}, \mathbf{N}_{\gamma\delta}^b\widehat{\mathbf{I}_{\delta\alpha}^b}^{-1} Y_{\alpha\delta}^b p_{\delta}^b] \\ &+ Y_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \mathbf{N}_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \mathcal{Y}_{\alpha\beta\gamma}^{ab}[I_{\beta\delta}p_{\delta}^{a-b}, I_{\gamma\epsilon}p_{\epsilon}^b] + Y_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \mathcal{C}_{\alpha\beta\gamma}^{ab}[\widehat{\mathbf{I}_{\beta\alpha}^{a-b}}^{-1} Y_{\alpha\beta}^{\pm a-b} p_{\beta}^{a-b}, I_{\gamma\epsilon}p_{\epsilon}^b] \\ &- \mathcal{A}_{\alpha\beta\gamma}^{ab}[\widehat{\mathbf{I}_{\beta\alpha}^{a-b}}^{-1} Y_{\alpha\beta}^{\pm a-b} p_{\beta}^{a-b}, \widehat{\mathbf{I}_{\gamma\alpha}^b}^{-1} Y_{\alpha\gamma}^{\pm b} p_{\gamma}^b] - \mathcal{B}_{\alpha\beta\gamma}^{ab}[I_{\beta\delta}p_{\delta}^{a-b}, I_{\gamma\epsilon}p_{\epsilon}^b] - \varepsilon_{\alpha\beta\gamma}^{ab}[\widehat{\mathbf{I}_{\beta\alpha}^{a-b}}^{-1} Y_{\alpha\beta}^{\pm a-b} p_{\beta}^{a-b}, I_{\gamma\epsilon}p_{\epsilon}^b] \\ &+ \mathcal{Y}_{\alpha\beta\gamma}^{ab}[I_{\beta\delta}p_{\delta}^{a-b}, \mathbf{H}_{\gamma\delta}^b p_{\delta}^b] + \mathcal{Y}_{\alpha\beta\gamma}^{ab}[\mathbf{H}_{\beta\delta}^{a-b} p_{\delta}^{a-b}, I_{\gamma\epsilon}p_{\epsilon}^b] + \mathbf{G}_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \mathcal{Y}_{\alpha\beta\gamma}^{ab}[I_{\beta\delta}p_{\delta}^{a-b}, I_{\gamma\epsilon}p_{\epsilon}^b] \\ &+ Y_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \mathcal{D}_{\alpha\beta\gamma}^{ab}[\widehat{\mathbf{I}_{\beta\alpha}^{a-b}}^{-1} Y_{\alpha\beta}^{\pm a-b} p_{\beta}^{a-b}, \widehat{\mathbf{I}_{\gamma\alpha}^b}^{-1} Y_{\alpha\gamma}^{\pm b} I_{\gamma\epsilon}p_{\epsilon}^b] = 0 \end{aligned} \quad (108)$$

$\Rightarrow$  quadratic-rank 5 tensor(2 upper, 3 lower):

$$\begin{aligned} &(\mathcal{Y}'_{\alpha\beta\gamma}^{ab}[I_{\beta\delta}, I_{\gamma\epsilon}]_{\alpha\beta\gamma}^{ab})_{\underline{\underline{p_{\delta}^{a-b}, p_{\epsilon}^b}}} + \mathcal{Y}_{\alpha\beta\gamma}^{ab}[\mathbf{N}_{\beta\delta}^{a-b}\widehat{\mathbf{I}_{\delta\alpha}^{a-b}}^{-1} Y_{\alpha\delta}^{a-b}, I_{\gamma\epsilon}]_{\underline{\underline{p_{\delta}^{a-b}, p_{\epsilon}^b}}} + \mathcal{Y}_{\alpha\beta\gamma}^{ab}[I_{\beta\delta}, \mathbf{N}_{\gamma\delta}^b\widehat{\mathbf{I}_{\delta\alpha}^b}^{-1} Y_{\alpha\delta}^b]_{\underline{\underline{p_{\delta}^{a-b}, p_{\epsilon}^b}}} \\ &+ Y_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \mathbf{N}_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \mathcal{Y}_{\alpha\beta\gamma}^{ab}[I_{\beta\delta}, I_{\gamma\epsilon}]_{\underline{\underline{p_{\delta}^{a-b}, p_{\epsilon}^b}}} + Y_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \mathcal{C}_{\alpha\beta\gamma}^{ab}[\widehat{\mathbf{I}_{\beta\alpha}^{a-b}}^{-1} Y_{\alpha\beta}^{\pm a-b}, I_{\gamma\epsilon}]_{\underline{\underline{p_{\delta}^{a-b}, p_{\epsilon}^b}}} \\ &- \mathcal{A}_{\alpha\beta\gamma}^{ab}[\widehat{\mathbf{I}_{\beta\alpha}^{a-b}}^{-1} Y_{\alpha\beta}^{\pm a-b}, \widehat{\mathbf{I}_{\gamma\alpha}^b}^{-1} Y_{\alpha\gamma}^{\pm b}]_{\underline{\underline{p_{\delta}^{a-b}, p_{\epsilon}^b}}} - \mathcal{B}_{\alpha\beta\gamma}^{ab}[I_{\beta\delta}, I_{\gamma\epsilon}]_{\underline{\underline{p_{\delta}^{a-b}, p_{\epsilon}^b}}} - \varepsilon_{\alpha\beta\gamma}^{ab}[\widehat{\mathbf{I}_{\beta\alpha}^{a-b}}^{-1} Y_{\alpha\beta}^{\pm a-b}, I_{\gamma\epsilon}]_{\underline{\underline{p_{\delta}^{a-b}, p_{\epsilon}^b}}} \\ &+ \mathcal{Y}_{\alpha\beta\gamma}^{ab}[I_{\beta\delta}, \mathbf{H}_{\gamma\delta}^b p_{\delta}^b] + \mathcal{Y}_{\alpha\beta\gamma}^{ab}[\mathbf{H}_{\beta\delta}^{a-b} p_{\delta}^{a-b}, I_{\gamma\epsilon}]_{\underline{\underline{p_{\delta}^{a-b}, p_{\epsilon}^b}}} + \mathbf{G}_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \mathcal{Y}_{\alpha\beta\gamma}^{ab}[I_{\beta\delta}, I_{\gamma\epsilon}]_{\underline{\underline{p_{\delta}^{a-b}, p_{\epsilon}^b}}} \\ &+ Y_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \mathcal{D}_{\alpha\beta\gamma}^{ab}[\widehat{\mathbf{I}_{\beta\alpha}^{a-b}}^{-1} Y_{\alpha\beta}^{\pm a-b}, \widehat{\mathbf{I}_{\gamma\alpha}^b}^{-1} Y_{\alpha\gamma}^{\pm b} I_{\gamma\epsilon}]_{\underline{\underline{p_{\delta}^{a-b}, p_{\epsilon}^b}}} = 0 \end{aligned} \quad (109)$$

These equation are solved from the outlet of the duct, applying the appropriate radiation boundary condition at the duct exit. Once  $Y(s)$  and  $\mathcal{Y}(s)$  are found through the duct, eq99 can then be used to replace the velocity modes with pressure modes in eq98, to obtain a numerically stable first

order ODE for the pressure modes:

$$\begin{aligned} \widehat{\mathbf{I}_{\alpha\beta}^a p'_\beta}^a &= \mathbf{N}_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} Y_{\alpha\beta}^a p_\beta^a + \mathbf{N}_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \mathcal{Y}_{\alpha\beta\gamma}^{ab} [p_\beta^{a-b}, p_\gamma^b] \\ &+ \mathbf{H}_{\alpha\beta}^a p_\beta^a + \mathcal{C}_{\alpha\beta\gamma}^{ab} [\widehat{\mathbf{I}_{\beta\alpha}^{a-b}}^{-1} Y_{\alpha\beta}^{\pm a-b} p_\beta^{a-b}, p_\gamma^b] + \mathcal{D}_{\alpha\beta\gamma}^{ab} [\widehat{\mathbf{I}_{\beta\alpha}^{a-b}}^{-1} Y_{\alpha\beta}^{\pm a-b} p_\beta^{a-b}, \widehat{\mathbf{I}_{\gamma\alpha}^b}^{-1} Y_{\alpha\gamma}^{\pm b} p_\gamma^b] \end{aligned} \quad (110)$$

This equation can be solved from the source to the outlet. As the equation involves the local admittance at each point, the solution includes both forward and backwards propagating waves together with their nonlinear interaction.

## VII. Boundary Conditions for an infinite uniform duct outlet

The simplest boundary condition to consider for the admittance is that of an outlet consisting of an infinitely long uniform duct of constant curvature for which we have only propagating waves and decaying evanescent waves. In such a duct no point can be distinguished from another longitudinally, therefore we must have the admittance being a fixed point of the admittance equations. To find the fixed points, we begin by combining eq99, ignoring the quadratic terms for the moment, to form a second order ODE for the pressure modes, G,H, the derivatives of M and N vanish:

$$\begin{aligned} \widehat{\mathbf{I}_{\alpha\beta}^a u'_\beta}^a + \mathbf{M}_{\alpha\beta}^a p_\beta^a &= 0 \\ \widehat{\mathbf{I}_{\alpha\beta}^a p'_\beta}^a - \mathbf{N}_{\alpha\beta}^a u_\beta^a &= 0 \end{aligned} \quad (111)$$

$$\widehat{\mathbf{I}_{\alpha\beta}^a p''_\beta}^a(s) + \mathbf{N}_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \mathbf{M}_{\alpha\beta}^a p_\beta^a(s) = 0 \quad (112)$$

$$\{v_i, -\lambda_i^2\} = \text{eig}(\mathbf{N}_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \mathbf{M}_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1}), \text{ suppose } \alpha = \beta \quad (113)$$

In matrix,

$$[\mathbf{N}_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \mathbf{M}_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1}]_{\alpha \times \alpha} V = V \Lambda^2, V = [v1, v2, v3, \dots], \Lambda = \text{diag}(i\lambda_1, i\lambda_2, \dots) \quad (114)$$

The solution in terms of forward and backward modes is given by:

$$p = p^+ + p^- = \sum_{k=1}^{\infty} (c_k^+ v_k e^{i\lambda_k s} + c_k^- v_k e^{-i\lambda_k s}) \quad (115)$$

where the  $v_i$  are the eigenvalue of NM with eigenvalues  $\lambda_i^2$ , with arbitrary  $c_k^+$  and  $c_k^-$ . Here, we have split the solution into forward and backward waves. The roots of the  $\lambda_k$  are chosen as follows:

$$\lambda_k = \begin{cases} (\lambda_k^2)^{1/2}, \lambda_k^2 > 0 \\ i(-\lambda_k^2)^{1/2}, \lambda_k^2 < 0 \end{cases} \quad (116)$$

to ensure either propagating or decaying evanescent modes in the positive direction. Based on extensive numerical evaluations, we observe that all of the eigenvectors of NM are real. We now introduce the characteristic forward and backwards admittance, linearly relating the forward and backwards modes:

$$\widehat{\mathbf{I}_{\alpha\beta}^a} u_{\beta}^{\pm a} = Y_{\alpha\beta}^{\pm a} p_{\beta}^{\pm a} \quad (117)$$

Using this, together with the linear equation relating pressure and velocity  $(p_{\beta}^{\pm a})' = N_{\alpha\beta}^a u_{\beta}^{\pm a}$ , we obtain an expression for  $Y_{\alpha\beta}^{\pm a}$

$$(p_{\beta}^{\pm a}(s))' = N_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} Y_{\alpha\beta}^{\pm a} p_{\beta}^{\pm a}(s) \quad (118)$$

Which is similar to above eig property, we have:

$$\begin{aligned} & [N_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} Y_{\alpha\beta}^{\pm a} \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1}]_{\alpha \times \alpha} V = \pm V \Lambda, V = [v1, v2, v3, \dots], \Lambda = \text{diag}(i\lambda_1, i\lambda_2, \dots) \\ \Rightarrow Y_{\alpha\beta}^{\pm a} &= \pm \widehat{\mathbf{I}_{\alpha\beta}^a} N_{\beta\alpha}^a -1 V \Lambda V^{-1} \widehat{\mathbf{I}_{\alpha\beta}^a} = \pm \widehat{\mathbf{I}_{\alpha\beta}^a} N_{\beta\alpha}^a -1 \sqrt{V \Lambda^2 V^{-1}} \widehat{\mathbf{I}_{\alpha\beta}^a} = \pm i \widehat{\mathbf{I}_{\alpha\beta}^a} N_{\beta\alpha}^a -1 \sqrt{\mathbf{N}_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \mathbf{M}_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \widehat{\mathbf{I}_{\alpha\beta}^a}} \end{aligned} \quad (119)$$

Substitute into eq, again ignoring G and H as the duct is assumed uniform:

$$\text{linear } -2D : Y_{\alpha\beta}'^a + Y_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \mathbf{N}_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} Y_{\alpha\beta}^a + \mathbf{M}_{\alpha\beta}^a + Y_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \mathbf{H}_{\alpha\beta}^a + \mathbf{G}_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} Y_{\alpha\beta}^a = 0, \quad (120)$$

$$Y_{\alpha\beta}'^a = \widehat{\mathbf{I}_{\alpha\beta}^a} N_{\beta\alpha}^a -1 \sqrt{\mathbf{N}_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \mathbf{M}_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \widehat{\mathbf{I}_{\alpha\beta}^a} \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \mathbf{N}_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \widehat{\mathbf{I}_{\alpha\beta}^a} N_{\beta\alpha}^a -1} \sqrt{\mathbf{N}_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \mathbf{M}_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \widehat{\mathbf{I}_{\alpha\beta}^a}} - M_{\alpha\beta}^a = 0 \quad (121)$$

Therefore,  $Y = Y^+$  is the boundary condition applied at exit, implying only outgoing (not ingoing) propagating waves and decaying evanescent waves in the outlet.

Now, we introduce a matrix  $\mathbf{W} = \mathbf{V}$  with columns given by the eigenvectors of  $Y^{\pm} N$  with corresponding eigenvalue matrix  $\pm \Lambda$ :

$$\begin{aligned} Y_{\alpha\beta}^{\pm a} &= \pm \widehat{\mathbf{I}_{\alpha\beta}^a} N_{\beta\alpha}^a -1 V_{\alpha \times \alpha}^a \Lambda_{\alpha \times \alpha}^a V_{\alpha \times \alpha}^a -1 \widehat{\mathbf{I}_{\alpha\beta}^a} \Rightarrow N_{\alpha\beta}^a \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} Y_{\alpha\beta}^{\pm a} \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} V_{\alpha \times \alpha}^a = \pm V_{\alpha \times \alpha}^a \Lambda_{\alpha \times \alpha}^a \\ Y_{\alpha\beta}^{\pm a} &= \pm \widehat{\mathbf{I}_{\alpha\beta}^a} W_{\beta \times \beta}^a \Lambda_{\beta \times \beta}^a W_{\beta \times \beta}^a -1 N_{\beta\alpha}^a -1 \widehat{\mathbf{I}_{\alpha\beta}^a} \Rightarrow \end{aligned} \quad (122)$$

We know that:

$$\begin{aligned}
(\mathcal{A}[x, y])_\alpha^a &= (\mathcal{A}_{\alpha\beta\gamma}^{ab}[x_\beta^{a-b}, y_\gamma^b])_\alpha^a = \sum_{b=-\infty}^{\infty} \sum_{\beta, \gamma=0}^{\infty} A_{\alpha\beta\gamma}^{ab} x_\beta^{a-b} y_\gamma^b \\
(\mathcal{A}[X, Y])_{\alpha\beta\gamma}^{ab} &= (\mathcal{A}_{\alpha\delta\epsilon}^{ab}[X_{\delta\beta}^{a-b}, Y_{\epsilon\gamma}^b])_{\alpha\beta\gamma}^{ab} = \sum_{\delta, \epsilon=0}^{\infty} A_{\alpha\delta\epsilon}^{ab} X_{\delta\beta}^{a-b} Y_{\epsilon\gamma}^b \\
\text{Thus, } \{(\mathcal{A}[X_1, Y_1])_{\alpha\beta\gamma}^{ab}[X_2, Y_2]\}_{\alpha\delta\epsilon}^{ab} &= ((\mathcal{A}_{\alpha\delta\epsilon}^{ab}[X_1_{\delta\beta}^{a-b}, Y_1_{\epsilon\gamma}^b])_{\alpha\beta\gamma}^{ab}[X_2_{\beta\delta}^{a-b}, Y_2_{\gamma\epsilon}^b])_{\alpha\delta\epsilon}^{ab} \\
&= \sum_{\beta, \gamma=0}^{\infty} \left( \sum_{\delta, \epsilon=0}^{\infty} A_{\alpha\delta\epsilon}^{ab} X_1_{\delta\beta}^{a-b} Y_1_{\epsilon\gamma}^b \right) X_2_{\beta\delta}^{a-b} Y_2_{\gamma\epsilon}^b \\
&= \sum_{\delta, \epsilon=0}^{\infty} \left\{ A_{\alpha\delta\epsilon}^{ab} \sum_{\beta=0}^{\infty} (X_1_{\delta\beta}^{a-b} X_2_{\beta\delta}^{a-b}) \sum_{\gamma=0}^{\infty} (Y_1_{\epsilon\gamma}^b Y_2_{\gamma\epsilon}^b) \right\} \\
&= (\mathcal{A}_{\alpha\delta\epsilon}^{ab}[X_1_{\delta\beta}^{a-b} X_2_{\beta\delta}^{a-b}, Y_1_{\epsilon\gamma}^b Y_2_{\gamma\epsilon}^b])_{\alpha\delta\epsilon}^{ab}
\end{aligned} \tag{123}$$

With  $\varepsilon = 0, G = 0, H = 0, \mathcal{Y}' = 0$ , fix points of the nonlinear admittance equation satisfy:

$$\begin{aligned}
&\mathcal{Y}_{\alpha\beta\gamma}^{ab} [\mathbf{N}_{\beta\delta}^{a-b} \widehat{\mathbf{I}}_{\delta\alpha}^{a-b}{}^{-1} Y_{\alpha\delta}^{a-b}, I_{\gamma\epsilon}] + \mathcal{Y}_{\alpha\beta\gamma}^{ab} [I_{\beta\delta}, \mathbf{N}_{\gamma\delta}^b \widehat{\mathbf{I}}_{\delta\alpha}^{b}{}^{-1} Y_{\alpha\delta}^b] \\
&+ Y_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^{a}{}^{-1} \mathbf{N}_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^{a}{}^{-1} \mathcal{Y}_{\alpha\beta\gamma}^{ab} [I_{\beta\delta}, I_{\gamma\epsilon}] + Y_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^{a}{}^{-1} \mathcal{C}_{\alpha\beta\gamma}^{ab} [\widehat{\mathbf{I}}_{\beta\alpha}^{a-b}{}^{-1} Y_{\alpha\beta}^{\pm a-b}, I_{\gamma\epsilon}] \\
&- \mathcal{A}_{\alpha\beta\gamma}^{ab} [\widehat{\mathbf{I}}_{\beta\alpha}^{a-b}{}^{-1} Y_{\alpha\beta}^{\pm a-b}, \widehat{\mathbf{I}}_{\gamma\alpha}^{b}{}^{-1} Y_{\alpha\gamma}^{\pm b}] - \mathcal{B}_{\alpha\beta\gamma}^{ab} [I_{\beta\delta}, I_{\gamma\epsilon}] = 0
\end{aligned} \tag{124}$$


We apply  $\{W_{\alpha\xi}^a\}^{-1}$  on the left of this equation and  $V_{\alpha\beta}^a$  to the right on both terms in the square brackets:

$$\begin{aligned}
&W^{-1} {}^a_{\xi\alpha} \mathcal{Y}_{\alpha\beta\gamma}^{\pm ab} [\mathbf{N}_{\beta\delta}^{a-b} \widehat{\mathbf{I}}_{\delta\alpha}^{a-b}{}^{-1} Y_{\alpha\delta}^{a-b} V, I_{\gamma\epsilon} V] + W^{-1} {}^a_{\xi\alpha} \mathcal{Y}_{\alpha\beta\gamma}^{\pm ab} [I_{\beta\delta} V, \mathbf{N}_{\gamma\delta}^b \widehat{\mathbf{I}}_{\delta\alpha}^{b}{}^{-1} Y_{\alpha\delta}^b V] \\
&+ W^{-1} {}^a_{\xi\alpha} Y_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^{a}{}^{-1} \mathbf{N}_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^{a}{}^{-1} \mathcal{Y}_{\alpha\beta\gamma}^{ab} [I_{\beta\delta} V, I_{\gamma\epsilon} V] + W^{-1} {}^a_{\xi\alpha} Y_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^{a}{}^{-1} \mathcal{C}_{\alpha\beta\gamma}^{ab} [\widehat{\mathbf{I}}_{\beta\alpha}^{a-b}{}^{-1} Y_{\alpha\beta}^{\pm a-b} V, I_{\gamma\epsilon} V] \\
&- W^{-1} {}^a_{\xi\alpha} \mathcal{A}_{\alpha\beta\gamma}^{ab} [\widehat{\mathbf{I}}_{\beta\alpha}^{a-b}{}^{-1} Y_{\alpha\beta}^{\pm a-b} V, \widehat{\mathbf{I}}_{\gamma\alpha}^{b}{}^{-1} Y_{\alpha\gamma}^{\pm b} V] - W^{-1} {}^a_{\xi\alpha} \mathcal{B}_{\alpha\beta\gamma}^{ab} [I_{\beta\delta} V, I_{\gamma\epsilon} V] = 0
\end{aligned} \tag{125}$$

$$\begin{aligned}
&W^{-1} {}^a_{\xi\alpha} \mathcal{Y}_{\alpha\beta\gamma}^{\pm ab} [\pm (V\Lambda)_{\beta\delta}^{a-b}, V_{\gamma\epsilon}^b] + W^{-1} {}^a_{\xi\alpha} \mathcal{Y}_{\alpha\beta\gamma}^{\pm ab} [V_{\beta\delta}^{a-b}, \pm (V\Lambda)_{\gamma\epsilon}^b] \\
&+ W^{-1} {}^a_{\xi\alpha} Y_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^{a}{}^{-1} \mathbf{N}_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^{a}{}^{-1} \mathcal{Y}_{\alpha\beta\gamma}^{ab} [V_{\beta\delta}^{a-b}, V_{\gamma\epsilon}^b] + W^{-1} {}^a_{\xi\alpha} Y_{\alpha\beta}^a \widehat{\mathbf{I}}_{\beta\alpha}^{a}{}^{-1} \mathcal{C}_{\alpha\beta\gamma}^{ab} [\widehat{\mathbf{I}}_{\beta\alpha}^{a-b}{}^{-1} Y_{\alpha\beta}^{\pm a-b} V_{\beta\delta}^{a-b}, V_{\gamma\epsilon}^b] \\
&- W^{-1} {}^a_{\xi\alpha} \mathcal{A}_{\alpha\beta\gamma}^{ab} [\widehat{\mathbf{I}}_{\beta\alpha}^{a-b}{}^{-1} Y_{\alpha\beta}^{\pm a-b} V_{\beta\delta}^{a-b}, \widehat{\mathbf{I}}_{\gamma\alpha}^{b}{}^{-1} Y_{\alpha\gamma}^{\pm b} V_{\gamma\epsilon}^b] - W^{-1} {}^a_{\xi\alpha} \mathcal{B}_{\alpha\beta\gamma}^{ab} [V_{\beta\delta}^{a-b}, I_{\gamma\epsilon} V_{\gamma\epsilon}^b] = 0
\end{aligned} \tag{126}$$

Next, we transform  $\mathcal{Y}^\pm$  in the following manner:

$$\mathcal{Y}^\pm[x, y] = W \tilde{\mathcal{Y}}^\pm[V^{-1}x, V^{-1}y] \tag{127}$$



**Fig. 1** 3D – model – lambda

$$\begin{aligned}
& \tilde{\mathcal{Y}}_{\alpha\beta\gamma}^{\pm ab}[\pm(\Lambda)_{\beta\delta}^{a-b}, I_{\gamma\epsilon}^b] + \tilde{\mathcal{Y}}_{\alpha\beta\gamma}^{\pm ab}[I_{\beta\delta}^{a-b}, \pm(\Lambda)_{\gamma\epsilon}^b] \pm \Lambda_{\xi\alpha}\tilde{\mathcal{Y}}_{\alpha\beta\gamma}^{\pm ab}[I_{\beta\delta}^{a-b}, I_{\gamma\epsilon}^b] \\
& + W^{-1}{}_{\xi\alpha}^a Y_{\alpha\beta}^{\pm a} \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \mathcal{C}_{\alpha\beta\gamma}^{ab} [\widehat{\mathbf{I}_{\beta\alpha}^{a-b}}^{-1} Y_{\alpha\beta}^{\pm a-b} V_{\beta\delta}^{a-b}, V_{\gamma\epsilon}^b] \quad (128) \\
& - W^{-1}{}_{\xi\alpha}^a \mathcal{A}_{\alpha\beta\gamma}^{ab} [\widehat{\mathbf{I}_{\beta\alpha}^{a-b}}^{-1} Y_{\alpha\beta}^{\pm a-b} V_{\beta\delta}^{a-b}, \widehat{\mathbf{I}_{\gamma\alpha}^b}^{-1} Y_{\alpha\gamma}^{\pm b} V_{\gamma\epsilon}^b] - W^{-1}{}_{\xi\alpha}^a \mathcal{B}_{\alpha\beta\gamma}^{ab} [V_{\beta\delta}^{a-b}, I_{\gamma\epsilon} V_{\gamma\epsilon}^b] = 0
\end{aligned}$$

$$\begin{aligned}
& \tilde{\mathcal{Y}}_{\alpha\beta\gamma}^{\pm ab} = \frac{1}{\pm i\lambda_\alpha^a \pm i\lambda_\beta^{a-b} \pm i\lambda_\gamma^b} (-W^{-1}{}_{\xi\alpha}^a Y_{\alpha\beta}^{\pm a} \widehat{\mathbf{I}_{\beta\alpha}^a}^{-1} \mathcal{C}_{\alpha\beta\gamma}^{ab} [\widehat{\mathbf{I}_{\beta\alpha}^{a-b}}^{-1} Y_{\alpha\beta}^{\pm a-b} V_{\beta\delta}^{a-b}, V_{\gamma\epsilon}^b]) \\
& + W^{-1}{}_{\xi\alpha}^a \mathcal{A}_{\alpha\beta\gamma}^{ab} [\widehat{\mathbf{I}_{\beta\alpha}^{a-b}}^{-1} Y_{\alpha\beta}^{\pm a-b} V_{\beta\delta}^{a-b}, \widehat{\mathbf{I}_{\gamma\alpha}^b}^{-1} Y_{\alpha\gamma}^{\pm b} V_{\gamma\epsilon}^b] + W^{-1}{}_{\xi\alpha}^a \mathcal{B}_{\alpha\beta\gamma}^{ab} [V_{\beta\delta}^{a-b}, I_{\gamma\epsilon} V_{\gamma\epsilon}^b])_{\alpha\beta\gamma}^{ab} \quad (129)
\end{aligned}$$

### VIII. Boundary Conditions for an infinite Helical duct outlet

We begin by considering the linear forms of equation eq98:

$$\begin{pmatrix} u \\ p \end{pmatrix}' = \begin{pmatrix} -G & -M \\ N & H \end{pmatrix} \begin{pmatrix} u \\ p \end{pmatrix} = L \begin{pmatrix} u \\ p \end{pmatrix} \quad (130)$$

$$\begin{pmatrix} u \\ p \end{pmatrix}'' = L^2 \begin{pmatrix} u \\ p \end{pmatrix} = \begin{pmatrix} -(G^2 - MN) & (GM - MH) \\ (HN - NG) & (H^2 - NM) \end{pmatrix} \begin{pmatrix} u \\ p \end{pmatrix} \quad (131)$$

This equation can be solved by  $u = \sum v_i^u e^{\lambda_i s}$  and  $p = \sum v_i^p e^{\lambda_i s}$ , similarly, we have:

$$\left\{ \begin{pmatrix} v_i^u \\ v_i^p \end{pmatrix}, \lambda_i \right\} = \text{eig}(L) \quad (132)$$

$$\left\{ \begin{pmatrix} v_i^u \\ v_i^p \end{pmatrix}, \lambda_i^2 \right\} = \text{eig}(L^2) \quad (133)$$

## IX. Open Ducts

### X. Separating the $\Psi$ intergrals into radial and angular parts

With  $\int_0^{2\pi} e^{im\phi} e^{-im\theta} d\theta = 2\pi$ ,  $\psi_{\alpha_{m\mu}}(r) = C_{\alpha_{m\mu}} J_m(\frac{j'_{m\mu} r}{h}) e^{im\phi}$ , we have:

$$C_{\alpha_{m\mu}} = \frac{(i)^m}{\sqrt{(\pi h^2(1 - \frac{m^2}{j'^2_{m\mu}})J_m^2(j'_{m\mu}))}} \quad (134)$$

except for :  $C_{\alpha_{01}} = \frac{1}{\sqrt{\pi}h}$

$$\psi_{\alpha_{m\mu}}(r) = C_{\alpha_{m\mu}} J_m(\frac{j'_{m\mu} r}{h}) e^{im\phi} \quad (135)$$

By normalization, h is not bother the equation of I.:

$$\begin{aligned} \int_0^h C1C2J1J2rdr &= \int_0^1 \frac{C_1^\#}{h} \frac{C_2^\#}{h} J1(j'_1 z) J2(j'_2 z) z h h dz = \int_0^1 C_1^\# C_2^\# 2J1(j'_1 z) J2(j'_2 z) z dz. \\ \int_0^h C1C2C3J1J2J3rdr &= \int_0^1 \frac{C_1^\#}{h} \frac{C_2^\#}{h} \frac{C_3^\#}{h} J1(j'_1 z) J2(j'_2 z) J3(j'_3 z) z h h dz = \\ \frac{1}{h} \int_0^1 C_1^\# C_2^\# C_3^\# 2J1(j'_1 z) J2(j'_2 z) J3(j'_3 z) z dz. \end{aligned}$$

Which this derivation could be used to simplify the code.

$$\begin{aligned} \Psi_{[\alpha](\beta)\gamma}[r(1 - \kappa r \cos\phi)] &= \int_0^{2\pi} \int_0^h [\frac{\partial \psi_\alpha}{\partial r}] [\frac{\partial \psi_\beta}{\partial \theta}] \psi_\gamma[r(1 - \kappa r \cos\phi)] dr d\theta \\ &= \mathcal{X}_{[\alpha]\beta\gamma}[r] \Theta_{\alpha(\beta)\gamma} - \kappa \mathcal{X}_{[\alpha]\beta\gamma}[r^2] \Theta_{\alpha(\beta)\gamma}[\cos\phi] \end{aligned} \quad (136)$$

with:

$$\begin{aligned} \mathcal{X}_{[\alpha]\beta\gamma} &= \int_0^h \frac{d}{dr} (C_{\alpha_{m\mu}} J_m(\frac{j'_{m\mu} r}{h})) C_{\beta_{n\nu}} J_n(\frac{j'_{\beta_{n\nu}} r}{h}) C_{\gamma_{k_w}} J_k(\frac{j'_{\gamma_{k_w}} r}{h}) dr \\ \Theta_{\alpha(\beta)\gamma} &= \int_0^{2\pi} e^{im\phi} \frac{d}{d\theta} (e^{in\phi}) e^{ik\phi} d\theta \end{aligned} \quad (137)$$

Bessel function recurrence relations are:

$$\begin{aligned} J_{m-1}(x) + J_{m+1}(x) &= 2m/x J_m(x) \\ J_{m-1}(x) - J_{m+1}(x) &= 2J'_m(x) \end{aligned} \quad (138)$$

we can have:

$$\begin{aligned} 2J_{m+1}(x) &= 2m/x J_m(x) - 2J'_m(x) \rightarrow \\ J'_m(x) &= m/x J_m(x) - J_{m+1}(x) \end{aligned} \quad (139)$$

$$\begin{aligned}
\mathcal{X}_{[\alpha]\beta}[r] &= \int_0^h \frac{d}{dr} (C_{\alpha_{m\mu}} J_m(\frac{j'_{\alpha_{m\mu}} r}{h})) C_{\beta_{nv}} J_n(\frac{j'_{\beta_{nv}} r}{h}) r dr \\
&= \int_0^h \frac{j'_{\alpha_{m\mu}}}{h} C_{\alpha_{m\mu}} [\frac{m}{j'_{\alpha_{m\mu}} r / h} J_m(\frac{j'_{\alpha_{m\mu}} r}{h}) - J_{m+1}(\frac{j'_{\alpha_{m\mu}} r}{h})] C_{\beta_{nv}} J_n(\frac{j'_{\beta_{nv}} r}{h}) r dr \\
&= C_{\alpha_{m\mu}} C_{\beta_{nv}} \int_0^h [m J_m(\frac{j'_{\alpha_{m\mu}} r}{h}) - \frac{j'_{\alpha_{m\mu}} r}{h} J_{m+1}(\frac{j'_{\alpha_{m\mu}} r}{h})] J_n(\frac{j'_{\beta_{nv}} r}{h}) dr
\end{aligned} \tag{140}$$

The  $\Theta$  intergrals can be calculated analytically:

$$\Theta_{\alpha\beta} = \int_0^{2\pi} e^{im\phi} e^{in\phi} d\theta = \begin{cases} 0 & \\ 2\pi, m+n=0 & \end{cases} = 2\pi\delta_{m,-n} \tag{141}$$

With Euler's equation:

$$\begin{aligned}
e^{ix} &= \cos x + i \sin x \\
\cos x &= [e^{ix} + e^{-ix}] / 2 \\
\sin x &= [e^{ix} - e^{-ix}] / 2i
\end{aligned} \tag{142}$$

$$\begin{aligned}
\Theta_{\alpha\beta}[\cos\phi] &= \int_0^{2\pi} \cos(\theta - \theta_0) e^{im\phi} e^{in\phi} d\theta = 1/2 \int_0^{2\pi} [e^{i(\theta-\theta_0)} + e^{-i(\theta-\theta_0)}] e^{im\phi} e^{in\phi} d\theta \\
&= 1/2 \int_0^{2\pi} e^{i(\theta-\theta_0)} e^{im\phi} e^{in\phi} d\theta + 1/2 \int_0^{2\pi} [e^{-i(\theta-\theta_0)}] e^{im\phi} e^{in\phi} d\theta \\
&= \pi e^{-i(1+m+n)\theta_0} \delta_{m,-n-1} + \pi e^{i(m+n-1)\theta_0} \delta_{m,-n+1} \\
&= \pi \delta_{m,-n-1} + \pi \delta_{m,-n+1}
\end{aligned} \tag{143}$$

$$\Theta_{(\alpha)\beta} = \int_0^{2\pi} \frac{\partial}{\partial\theta} (e^{im\phi}) e^{in\phi} d\theta = 2\pi im \delta_{m,-n} \tag{144}$$

$$\Theta_{(\alpha)\beta}[\cos\phi] = im[\pi\delta_{m,-n-1} + \pi\delta_{m,-n+1}] \tag{145}$$

$$\Theta_{\alpha\beta\gamma} = \int_0^{2\pi} e^{im\phi} e^{in\phi} e^{ik\theta} d\theta = 2\pi \delta_{m,-n,-k} \tag{146}$$

$$\Theta_{\alpha\beta\gamma}[\cos\phi] = 1/2 \int_0^{2\pi} [e^{i(\theta-\theta_0)} + e^{-i(\theta-\theta_0)}] e^{im\phi} e^{in\phi} e^{ik\theta} d\theta = \pi\delta_{m,-n,-k-1} + \pi\delta_{m,-n,-k+1} \tag{147}$$

$$\Theta_{\alpha\beta\gamma}[\sin\phi] = 1/(2i) \int_0^{2\pi} [e^{i(\theta-\theta_0)} - e^{-i(\theta-\theta_0)}] e^{im\phi} e^{in\phi} e^{ik\theta} d\theta = -i[\pi\delta_{m,-n,-k-1} - \pi\delta_{m,-n,-k+1}]$$

$$(148)$$

$$\Theta_{(\alpha)\beta\gamma} = im \int_0^{2\pi} e^{im\phi} e^{in\phi} e^{ik\phi} d\theta = 2\pi im\delta_{m,-n,-k}$$

$$(149)$$

$$\Theta_{(\alpha)\beta\gamma}[\cos\phi] = 1/2 \int_0^{2\pi} [e^{i(\theta-\theta_0)} + e^{-i(\theta-\theta_0)}] e^{im\phi} e^{in\phi} e^{ik\theta} d\theta = im[\pi\delta_{m,-n,-k-1} + \pi\delta_{m,-n,-k+1}]$$

$$(150)$$

We also have:

$$\Psi_{\{\alpha\}\beta} = \mathcal{X}_{\{\alpha\}\beta} \Theta_{\alpha\beta} + \mathcal{X}_{\alpha\beta} \Theta_{\{\alpha\}\beta} \quad (151)$$

with:

$$\begin{aligned} \mathcal{X}_{\{\alpha\}\beta} &= \int_0^h \frac{d}{ds} (C_{\alpha_{m\mu}}(s) J_m(\frac{j'_{\alpha_{m\mu}} r}{h(s)})) C_{\beta_{n\nu}} J_n(\frac{j'_{\beta_{n\nu}} r}{h}) dr \\ &= \int_0^h [(\frac{dC_{\alpha_{m\mu}}(s)}{ds}) J_m(\frac{j'_{\alpha_{m\mu}} r}{h(s)}) + C_{\alpha_{m\mu}} J_m(\frac{j'_{\alpha_{m\mu}} r}{h(s)}) \frac{-j'_{\alpha_{m\mu}} r}{h(s)^2} \frac{dh(s)}{ds}] C_{\beta_{n\nu}} J_n(\frac{j'_{\beta_{n\nu}} r}{h}) dr \\ &= \int_0^h [(\frac{d}{ds} \frac{1}{h(s)}) C_{\alpha_{m\mu}}(s) J_m(\frac{j'_{\alpha_{m\mu}} r}{h(s)}) + C_{\alpha_{m\mu}} J_m(\frac{j'_{\alpha_{m\mu}} r}{h(s)}) \frac{-j'_{\alpha_{m\mu}} r}{h(s)^2} \frac{dh(s)}{ds}] C_{\beta_{n\nu}} J_n(\frac{j'_{\beta_{n\nu}} r}{h}) dr \\ &= \int_0^h [(-\frac{h'(s)}{h}) C_{\alpha_{m\mu}}(s) J_m(\frac{j'_{\alpha_{m\mu}} r}{h(s)}) - C_{\alpha_{m\mu}} \frac{\partial}{\partial r} (J_m(\frac{j'_{\alpha_{m\mu}} r}{h(s)})) \frac{h'(s)}{h} r] C_{\beta_{n\nu}} J_n(\frac{j'_{\beta_{n\nu}} r}{h}) dr \\ &\quad = -\frac{h'}{h} (\mathcal{X}_{[\alpha]\beta}[r] + \mathcal{X}_{\alpha\beta}) \end{aligned} \quad (152)$$

$$\begin{aligned} \mathcal{X}_{\{\alpha\}\beta}[r] &= \int_0^h \frac{d}{ds} (C_{\alpha_{m\mu}}(s) J_m(\frac{j'_{\alpha_{m\mu}} r}{h(s)})) C_{\beta_{n\nu}} J_n(\frac{j'_{\beta_{n\nu}} r}{h}) r dr \\ &= \int_0^h [(\frac{dC_{\alpha_{m\mu}}(s)}{ds}) J_m(\frac{j'_{\alpha_{m\mu}} r}{h(s)}) + C_{\alpha_{m\mu}} J_m(\frac{j'_{\alpha_{m\mu}} r}{h(s)}) \frac{-j'_{\alpha_{m\mu}} r}{h(s)^2} \frac{dh(s)}{ds}] C_{\beta_{n\nu}} J_n(\frac{j'_{\beta_{n\nu}} r}{h}) r dr \\ &= \int_0^h [(\frac{d}{ds} \frac{1}{h(s)}) C_{\alpha_{m\mu}}(s) J_m(\frac{j'_{\alpha_{m\mu}} r}{h(s)}) + C_{\alpha_{m\mu}} J_m(\frac{j'_{\alpha_{m\mu}} r}{h(s)}) \frac{-j'_{\alpha_{m\mu}} r}{h(s)^2} \frac{dh(s)}{ds}] C_{\beta_{n\nu}} J_n(\frac{j'_{\beta_{n\nu}} r}{h}) r dr \\ &= \int_0^h [(-\frac{h'(s)}{h}) C_{\alpha_{m\mu}}(s) J_m(\frac{j'_{\alpha_{m\mu}} r}{h(s)}) - C_{\alpha_{m\mu}} \frac{\partial}{\partial r} (J_m(\frac{j'_{\alpha_{m\mu}} r}{h(s)})) \frac{h'(s)}{h} r] C_{\beta_{n\nu}} J_n(\frac{j'_{\beta_{n\nu}} r}{h}) r dr \\ &\quad = -\frac{h'}{h} (\mathcal{X}_{[\alpha]\beta}[r^2] + \mathcal{X}_{\alpha\beta}[r]) \end{aligned} \quad (153)$$

$$\begin{aligned}
\mathcal{X}_{\{\alpha\}\beta\gamma} &= \int_0^h \frac{d}{ds} (C_{\alpha_{m\mu}}(s) J_m(\frac{j'_{\alpha_{m\mu}} r}{h(s)})) C_{\beta_{nv}} J_n(\frac{j'_{\beta_{nv}} r}{h}) C_{\gamma_{kw}} J_n(\frac{j'_{\gamma_{kw}} r}{h}) dr \\
&= \int_0^h [(\frac{dC_{\alpha_{m\mu}}(s)}{ds}) J_m(\frac{j'_{\alpha_{m\mu}} r}{h(s)}) + C_{\alpha_{m\mu}} J_m(\frac{j'_{\alpha_{m\mu}} r}{h(s)}) \frac{-j'_{\alpha_{m\mu}} r}{h(s)^2} \frac{dh(s)}{ds}] C_{\beta_{nv}} J_n(\frac{j'_{\beta_{nv}} r}{h}) C_{\gamma_{kw}} J_n(\frac{j'_{\gamma_{kw}} r}{h}) dr \\
&= \int_0^h [(\frac{d\frac{1}{h(s)}}{ds}) C_{\alpha_{m\mu}}(s) J_m(\frac{j'_{\alpha_{m\mu}} r}{h(s)}) + C_{\alpha_{m\mu}} J_m(\frac{j'_{\alpha_{m\mu}} r}{h(s)}) \frac{-j'_{\alpha_{m\mu}} r}{h(s)^2} \frac{dh(s)}{ds}] C_{\beta_{nv}} J_n(\frac{j'_{\beta_{nv}} r}{h}) C_{\gamma_{kw}} J_n(\frac{j'_{\gamma_{kw}} r}{h}) dr \\
&= \int_0^h [(-\frac{h'(s)}{h}) C_{\alpha_{m\mu}}(s) J_m(\frac{j'_{\alpha_{m\mu}} r}{h(s)}) - C_{\alpha_{m\mu}} \frac{\partial}{\partial r} (J_m(\frac{j'_{\alpha_{m\mu}} r}{h(s)})) \frac{h'(s)}{h} r] C_{\beta_{nv}} J_n(\frac{j'_{\beta_{nv}} r}{h}) C_{\gamma_{kw}} J_n(\frac{j'_{\gamma_{kw}} r}{h}) dr \\
&\quad = -\frac{h'}{h} (\mathcal{X}_{[\alpha]\beta\gamma}[r] + \mathcal{X}_{\alpha\beta\gamma})
\end{aligned} \tag{154}$$

$$\begin{aligned}
\Theta_{\{\alpha\}\beta} &= \int_0^{2\pi} \frac{\partial}{\partial s} (e^{im\phi}) e^{in\phi} d\theta = -im \int_0^{2\pi} \frac{\partial \theta_0(s)}{\partial s} e^{im\phi} e^{in\phi} d\theta \\
&\quad = -\tau \Theta_{(\alpha)\beta} = -\tau 2\pi im \delta_{m,-n}
\end{aligned} \tag{155}$$

$$\Theta_{\alpha\{\beta\}} = -\tau \Theta_{\alpha(\beta)} = -\tau 2\pi in \delta_{m,-n} \tag{156}$$


Similarly,

$$\frac{d}{ds} (\Psi_{\alpha\beta\gamma}[r]) = \frac{d}{ds} (\mathcal{X}_{\alpha\beta\gamma}[r]) \Theta_{\alpha\beta\gamma} \tag{157}$$


$$\frac{d}{ds} \left( \int_0^{h(s)} f(r, s) dr \right) = \int_0^{h(s)} \frac{\partial}{\partial \alpha} f(r, s) dr + \frac{dh(s)}{ds} f(h(s), s) \tag{158}$$

with


$$\begin{aligned}
\frac{d}{ds} (\mathcal{X}_{\alpha\beta\gamma}[r]) &= \mathcal{X}_{\{\alpha\}\beta\gamma}[r] + \mathcal{X}_{\alpha\{\beta\}\gamma}[r] + \mathcal{X}_{\alpha\beta\{\gamma\}}[r] \\
&\quad + h' h [C_{\alpha_{m\mu}}(s) J_m(\frac{j'_{\alpha_{m\mu}} r}{h(s)}) C_{\beta_{nv}} J_n(\frac{j'_{\beta_{nv}} r}{h}) C_{\gamma_{kw}} J_n(\frac{j'_{\gamma_{kw}} r}{h})]^h \\
&= -\frac{h'}{h} (\mathcal{X}_{[\alpha]\beta\gamma}[r^2]) - \frac{h'}{h} (\mathcal{X}_{\alpha[\beta]\gamma}[r^2]) - \frac{h'}{h} (\mathcal{X}_{\alpha\beta[\gamma]}[r^2]) - 3 \frac{h'}{h} \mathcal{X}_{\alpha\beta\gamma}[r] \\
&\quad + h' h C_{\alpha_{m\mu}}(s) J_m(j'_{\alpha_{m\mu}}) C_{\beta_{nv}} J_n(j'_{\beta_{nv}}) C_{\gamma_{kw}} J_n(j'_{\gamma_{kw}})
\end{aligned} \tag{159}$$



**Fig. 2**  $\Theta$



**Fig. 3**  $X_\Theta$



**Fig. 4**  $\Psi$

- A. Example for matlab simulation
- B. Example for matlab simulation
- C. Example for matlab simulation

$$\begin{aligned}
Fig1 : \Psi_{\alpha\beta}[r] &= \mathcal{X}_{\alpha\beta}[r]\Theta_{\alpha\beta} := \int_0^h rC_{\alpha_{mu}}(s)J_m(\frac{j'_{\alpha_{mu}}r}{h})C_{\beta_{nv}}(scc)J_n(\frac{j'_{\beta_{nv}}r}{h})dr(2\pi\delta_{m,-n}) \\
Fig2 : \Psi_{\alpha\beta}[rcos\phi] &= \mathcal{X}_{\alpha\beta}[r]\Theta_{\alpha\beta}[cos\phi] := \int_0^h rC_{\alpha_{mu}}J_m(\frac{j'_{\alpha_{mu}}r}{h})C_{\beta_{nv}}J_n(\frac{j'_{\beta_{nv}}r}{h})dr(\pi\delta_{m,-n-1} + \pi\delta_{m,-n+1}) \\
Fig3 : \Psi_{\alpha\beta}[r^2cos\phi] &= \mathcal{X}_{\alpha\beta}[r^2]\Theta_{\alpha\beta}[cos\phi] := \int_0^h r^2C_{\alpha_{mu}}J_m(\frac{j'_{\alpha_{mu}}r}{h})C_{\beta_{nv}}J_n(\frac{j'_{\beta_{nv}}r}{h})dr(\pi\delta_{m,-n-1} + \pi\delta_{m,-n+1}) \\
Fig4 : \Psi_{\alpha\beta}[r^2sins\phi] &= \mathcal{X}_{\alpha\beta}[r^2]\Theta_{\alpha\beta}[sin\phi] := \int_0^h r^2C_{\alpha_{mu}}J_m(\frac{j'_{\alpha_{mu}}r}{h})C_{\beta_{nv}}J_n(\frac{j'_{\beta_{nv}}r}{h})dr(-i(\pi\delta_{m,-n-1} - \pi\delta_{m,-n+1})) \\
Fig5 : \Psi_{(\alpha)\beta}[r] &= \mathcal{X}_{\alpha\beta}[r]\Theta_{(\alpha)\beta} := \int_0^h rC_{\alpha_{mu}}J_m(\frac{j'_{\alpha_{mu}}r}{h})C_{\beta_{nv}}J_n(\frac{j'_{\beta_{nv}}r}{h})dr(2\pi im\delta_{m,-n}) \\
Fig6 : \Psi_{\{\alpha\}\beta} &= \mathcal{X}_{\{\alpha\}\beta}\Theta_{\alpha\beta} + \mathcal{X}_{\alpha\beta}\Theta_{\{\alpha\}\beta} \\
&\quad \mathcal{X}_{\{\alpha\}\beta} := -\frac{h'}{h}(\mathcal{X}_{[\alpha]\beta}[r] + \mathcal{X}_{\alpha\beta}) \\
&\quad \mathcal{X}_{[\alpha]\beta}[r] := \int_0^h r\frac{d}{dr}(C_{\alpha_{mu}}J_m(\frac{j'_{\alpha_{mu}}r}{h}))C_{\beta_{nv}}J_n(\frac{j'_{\beta_{nv}}r}{h})dr \\
&\quad \Theta_{\{\alpha\}\beta} = \int_0^{2\pi} \frac{\partial}{\partial s}(e^{im\phi})e^{in\phi}d\theta = -\tau 2\pi im\delta_{m,-n}
\end{aligned} \tag{160}$$

## XI. Tensors in matlab for numerical simulation

### A. Tensor times vectors: $\mathcal{A}\bar{\times}_nu$

Let  $\mathcal{A}$  be a tensor of size  $I_1 \times I_2 \times \dots \times I_N$ ,  $u$  be a vector of size  $I_n$ .

We have:

$$\begin{aligned}
ttv(\mathcal{A}, \{u\}, [n]) &= (\mathcal{A}\bar{\times}_nu)(i_1, \dots, i_{n-1}, i_{n+1}, \dots, i_N) \\
&\quad \sum_{i_n=1}^{I_n} \mathcal{A}(i_1, i_2, \dots, i_N)u(i_n)
\end{aligned} \tag{161}$$

$$\begin{aligned}
ttv(A_{m \times n}, \{u_{m \times 1}\}, [1]) &= A_{m \times n}\bar{\times}_1 u_{m \times 1} = A_{m \times n}^T u_{m \times 1}
\end{aligned} \tag{162}$$

$$ttv(A_{m \times n}, \{v_{n \times 1}\}, [2]) = A_{m \times n}\bar{\times}_2 v_{n \times 1} = A_{m \times n} v_{n \times 1}$$

Property:

$$\begin{aligned}
ttv(\mathcal{A}, \{u, v\}, [m, n]) &= \mathcal{A} \bar{\times}_m u \bar{\times}_n v \\
&= ttv(ttv(\mathcal{A}, \{u\}, [m]), \{v\}, [n - 1]) = (\mathcal{A} \bar{\times}_m u) \bar{\times}_{n-1} v \\
&= ttv(ttv(\mathcal{A}, \{v\}, [v]), \{u\}, [m]) = (\mathcal{A} \bar{\times}_n v) \bar{\times}_m u
\end{aligned} \tag{163}$$

Multiplication with a sequence of vectors

$$\beta = \mathcal{A} \bar{\times}_1 u^{(1)} \bar{\times}_2 u^{(2)} \dots \bar{\times}_N u^{(N)} = \mathcal{A} \bar{\times} u \tag{164}$$

*like* :  $ttv(X, \{A, B, C, D\}) = ttv(X, \{A, B, C, D\}, [1234]) = ttv(X, \{D, C, B, A\}, [4321])$

Multiplication with **all but one** of a sequence of vectors

$$\begin{aligned}
b &= \mathcal{A} \bar{\times}_1 u^{(1)} \bar{\times}_2 u^{(2)} \dots \bar{\times}_{n-1} u^{(2)} \bar{\times}_{n+1} u^{(2)} \dots \bar{\times}_N u^{(N)} = \mathcal{A} \bar{\times}_{-n} u \\
&\text{like : } X = tenrand([5, 3, 4, 2]);
\end{aligned} \tag{165}$$

$A = rand(5, 1); B = rand(3, 1); C = rand(4, 1); D = rand(2, 1);$

$$Y = ttv(X, \{A, B, D\}, -3) = ttv(X, \{A, B, C, D\}, -3)$$

## B. Tensor times matrix (ttm): $\mathcal{A} \times_n u$

Let  $\mathcal{A}$  be a tensor of size  $I_1 \times I_2 \times \dots \times I_N$ , U be a matrix of size  $J_n \times I_n$ .

We have:

$$\begin{aligned}
ttm(\mathcal{A}, \{U\}, [n]) &= (\mathcal{A} \times_n U)(i_1, \dots, i_{n-1}, j_n, i_{n+1}, \dots, i_N) \\
&= \sum_{i_n=1}^{I_n} \mathcal{A}(i_1, i_2, \dots, i_N) U(j_n, i_n)
\end{aligned} \tag{166}$$

*like* :  $X = tensor(rand(5, 3, 4, 2)); A = rand(4, 5);$

$$Y = ttm(X, A, 1) = ttm(X, \{A, B, C, D\}, 1) = ttm(X, A', 1, 't')$$

Matrix Interpretation

$$\begin{aligned}
ttm(A_{m \times n}, \{U_{m \times k}^T\}, [1]) &= A \times_1 U^T = U^T A \\
ttm(A_{m \times n}, \{V_{m \times k}^T\}, [2]) &= A \times_2 V^T = A V \\
ttm(A, \{U, V\}, [1, 2]) &= U A V^T
\end{aligned} \tag{167}$$

$Y = \text{ttm}(X, A, B, C, D, [1234]); \% <-- 4-way multiply.$

$Y = \text{ttm}(X, D, C, B, A, [4321]); \% <-- \text{Same as above}.$

$Y = \text{ttm}(X, A, B, C, D); \% <-- \text{Same as above}.$

$Y = \text{ttm}(X, A', B', C', D', t') \% <-- \text{Same as above}.$

$$Y = \text{ttm}(X, C, D, [34]); \% <-- X \times C \text{ in mode } -3 \text{ and } D \text{ in mode } -4 \quad (168)$$

$Y = \text{ttm}(X, A, B, C, D, [34]) \% <-- \text{Same as above}.$

$Y = \text{ttm}(X, A, B, D, [124]); \% <-- 3-way multiply.$

$Y = \text{ttm}(X, A, B, C, D, [124]); \% <-- \text{Same as above}.$

$Y = \text{ttm}(X, A, B, D, -3); \% <-- \text{Same as above}.$

$Y = \text{ttm}(X, A, B, C, D, -3) \% <-- \text{Same as above}.$

Property

$$\begin{aligned} \text{ttm}(\mathcal{A}, \{u, v\}, [m, n]) &= \mathcal{A} \times_m u \bar{\times}_n v \\ &= \text{ttm}(\text{ttm}(\mathcal{A}, \{u\}, [m]), \{v\}, [n]) = (\mathcal{A} \times_m u) \times_n v \\ &= \text{ttm}(\text{ttm}(\mathcal{A}, \{v\}, [v]), \{u\}, [m]) = (\mathcal{A} \times_n v) \times_m u \end{aligned} \quad (169)$$

### C. Tensor times tensor (ttt): $\langle \mathcal{A}, \mathcal{B} \rangle$

Let  $\mathcal{A}$  and  $\mathcal{B}$  be a tensor of size  $I_1 \times I_2 \times \dots \times I_N$ .

$$\begin{aligned} \langle \mathcal{A}, \mathcal{B} \rangle &= \\ \beta &= \sum_{i_1=1}^{I_1} \sum_{i_2=1}^{I_2} \dots \sum_{i_N=1}^{I_N} \mathcal{A}(i_1, i_2, \dots, i_N) \mathcal{B}(i_1, i_2, \dots, i_N) \end{aligned} \quad (170)$$

$X = \text{tensor}(\text{rand}(4, 2, 3)); Y = \text{tensor}(\text{rand}(3, 4, 2));$

$Z = \text{ttt}(X, Y); \% \text{ -- Outerproduct of } X \text{ and } Y.$

$\text{size}(Z)$

$Z = \text{ttt}(X, X, 1 : 3); \% \text{ -- Innerproduct of } X \text{ with itself.}$

(171)

$Z = \text{ttt}(X, Y, [123], [231]); \% \text{ -- Innerproduct of } XY.$

$Z = \text{innerprod}(X, \text{permute}(Y, [231])); \% \text{ -- Same as above.}$

$Z = \text{ttt}(X, Y, [13], [21]); \% \text{ -- Product of } XY \text{ along specified dims.}$

## XII. model of helical duct

### A. w

The duct is described by its centreline  $\mathbf{q}(s)$  at arclength  $s$  from the inlet of the duct adn the radial distance from the centreline  $h = h(s)$ . The general position vector ( $x$ ) in the duct is given in terms of  $(s, r, \theta)$ :

$$\mathbf{x} = \mathbf{q}(s) + r\cos(\theta - \theta_0)\hat{\mathbf{n}} + r\sin(\theta - \theta_0)\hat{\mathbf{b}} \quad (172)$$

where  $\hat{\mathbf{n}} = \hat{\mathbf{n}}(s)$  is the normal to the centreline and  $\hat{\mathbf{b}} = \hat{\mathbf{b}}(s)$  id the binormal given by  $\hat{\mathbf{b}} = \hat{\mathbf{t}} \times \hat{\mathbf{n}}$  for the tangent to the centreline  $\hat{\mathbf{t}} = \hat{\mathbf{t}}(s)$ . The vector  $\hat{\mathbf{n}}, \hat{\mathbf{b}}$  and  $\hat{\mathbf{t}}$  are related by the Frenet-Serret formulas:

$$\frac{d\hat{\mathbf{q}}}{ds} = \hat{\mathbf{t}}, \quad \frac{d\hat{\mathbf{t}}}{ds} = \kappa\hat{\mathbf{n}}, \quad \frac{d\hat{\mathbf{n}}}{ds} = -\kappa\hat{\mathbf{t}} + \tau\hat{\mathbf{b}}, \quad \frac{d\hat{\mathbf{b}}}{ds} = -\tau\hat{\mathbf{n}} \quad (173)$$

where  $\kappa = \kappa(s)$  is the local curvature of the duct and  $\tau = \text{tau}(s)$  is the torsion. Here, intorduce

$\theta'_0 = \tau$ , the cross-term differentials vanish and the metric reduces:

$$\begin{aligned}
d\mathbf{x} &= d(\mathbf{q}(s)) + d(r\cos(\theta - \theta_0)\hat{\mathbf{n}}) + d(r\sin(\theta - \theta_0)\hat{\mathbf{b}}) \\
&= \hat{\mathbf{t}}ds + dr\cos\phi\hat{\mathbf{n}} - r\hat{\mathbf{n}}\sin\phi(d\theta - \tau ds) + r\cos\phi(-\kappa\hat{\mathbf{t}} + \tau\hat{\mathbf{b}})ds \\
&\quad + dr\sin\phi\hat{\mathbf{b}} + r\hat{\mathbf{b}}\cos\phi(d\theta - \tau ds) + r\sin\phi(-\tau\hat{\mathbf{n}})ds \\
&= \hat{\mathbf{t}}(1 - \kappa r\cos\phi)ds + \hat{\mathbf{n}}(dr\cos\phi - r\sin\phi d\theta) + \hat{\mathbf{b}}(dr\sin\phi + r\cos\phi d\theta)
\end{aligned} \tag{174}$$

Thus,

$$\begin{aligned}
d\mathbf{x} \cdot d\mathbf{x} &= (1 - \kappa r\cos\phi)^2 ds^2 + (dr\cos\phi - r\sin\phi d\theta)^2 + (dr\sin\phi + r\cos\phi d\theta)^2 \\
&= (1 - \kappa r\cos\phi)^2 ds^2 + dr^2 + r^2 d\theta^2
\end{aligned} \tag{175}$$

As a result, we have an orthogonal coordinate system and as such do not need to distinguish between covariant and contravariant bases.

### XIII. funm-Evaluate general matrix function

<https://ww2.mathworks.cn/help/matlab/ref/funm.html>

### XIV. mtimesx-does a matrix multiply of two inputs

mtimesx is a fast general purpose matrix and scalar multiply routine that utilizes BLAS calls and custom code to perform the calculations. mtimesx also has extended support for n-Dimensional ( $nD$ ,  $n > 2$ ) arrays, treating these as arrays of 2D matrices for the purposes of matrix operations.

"Doesn't MATLAB already do this?" For 2D matrices, yes, it does. However, MATLAB does not always implement the most efficient algorithms for memory access, and MATLAB does not always take full advantage of symmetric cases. The mtimesx 'SPEED' mode attempts to do both of these to the fullest extent possible. For  $nD$  matrices, MATLAB does not have direct support for this. One is forced to write loops to accomplish the same thing that mtimesx can do faster.

Examples:

$$\begin{aligned}
C &= \text{mtimesx}(A, B) \% \text{performs the calculation} C = A * B \\
C &= \text{mtimesx}(A, T, B) \% \text{performs the calculation} C = A.' * B \\
C &= \text{mtimesx}(A, B, G) \% \text{performs the calculation} C = A * \text{conj}(B) \\
C &= \text{mtimesx}(A, C, B, C') \% \text{performs the calculation} C = A' * B'
\end{aligned} \tag{176}$$